

HYPERBOLIC VIBES AND PLASMONS-PHOTON CATAL-
PROMISSION TO RIGID METACRYLIC SYSTEMS

By

DAVID E. KYLE

A DISSERTATION PRESENTED TO THE GRADUATE COUNCIL OF
THE UNIVERSITY OF FLORIDA
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE
DEGREE OF DOCTOR OF PHILOSOPHY

UNIVERSITY OF FLORIDA

1976

in my presence
Margaret Hyltonier
and
Stephanie Anne Neta

ACKNOWLEDGMENTS

The author wishes to express his appreciation to his research director, Professor James H. Bayne, for suggesting this research problem and for the guidance and criticism received during the course of this investigation. The author also wishes to thank Mrs. Marie A. Binkley and Roy E. King for their counsel, advice, and contributions to his education at the University of Florida.

Special thanks are extended to Henry Maglich, Emmet McNeill, Jim Buchanan, Bill Baker, and the other members of his research group for their friendship, assistance and helpful suggestions.

The author is indebted to the Shanon Bookkeeper for the patience, efficiency and the sacrifices she made during the typing of this manuscript.

Finally, it is a pleasure to acknowledge the encouragement, companionship and efficiency so generously given of Linda Joanne Stone during the difficult periods leading to the final preparation of this dissertation.

TABLE OF CONTENTS

	Page
ACKNOWLEDGMENTS	iii
LIST OF TABLES	iv
LIST OF FIGURES	v
ABSTRACT	viii
CHAPTER I - INTRODUCTION (LIFE: EXPERIENCE AND INFORMATION OF 21-4-BUTYLENE-2-ETHYL-2-ETHYLENEFLUOROPOLYMER . . .	1
Introduction	1
Basic	21
Discussion	21
CHAPTER II - INTRODUCTION FLUOROPOLYMER SALTS: OLIGOCHLORIDE OF 21-4-BUTYLENE-2-ETHYL-2-ETHYLENEFLUOROPOLYMER . . .	24
Introduction	24
Basic	34
Discussion	34
CONCLUSION	61
Introduction	61
Basic	62
DISCUSSION	70
CHAPTER III - ANALYSIS OF 2-BUTYLENEFLUORINE TO ANALYSIS . .	74
Basic	74
Discussion	80
CHAPTER IV - EXPERIMENTAL	81
1-ethylfluorine Chloride (20)	84
2,3,4,5-tetraethyl-2-fluorine (20)	84
2,3,4,5-tetraethyl Chloride (24)	84

1,1,3-Trichloro-2-isopropyl-2-(<i>p</i> -Tolyl)-4-isothiazolylidene- 1,2,3,4-Tetrachloroquinazolin-1-one	1953
Acid Hydrolysis of 1,4-Bisacetyl-1-isothiazyl-3-(<i>p</i> -Tolyl)-4- isothiazolylidene-2,1,3,4-Tetrachloroquinazolin-1-one	1953
Isomethyl Chloride (158)	1953
2-Mercaptoisothiazolidine (90)	1953
Hydrolytic Action of Phosphoric Acid (271)	1953
1,1,3-Trichloro-2-(3,4,4-Trichloropropylideneazepidin-2-ylidene- 1-one)	1954
Acid Catalyzed Cyclization of Dimethylketone-8-Phenyl-1-one to 1,1,3-Trichloro-2-(3,4,4-Trichloropropylideneazepidin-2-ylidene- 1-one)	1954
1,1,3-Trichloro-2-(Isopropyl-4,4-Dimethyl-2-oxopropylidene-3,4,3,4- Tetrahydro- <i>n</i> -Triazin-1-one) Fluorination (181)	1955
Fluorination of 1,1,1-Trichloro-2-(3,4-Trichloropropylideneazepidin-2- ylidene-1-one) with SilverNitrate Acid	1955
Alkaline Hydrolysis of 1,1,3-Trichloro-2-(Isopropyl-4,4- Dimethylketone-3,4,3,4-Tetrahydro- <i>n</i> -Triazin-1-one) Fluorina- tion (185)	1955
1,1,3-Trichloro-2-(Isopropyl-4,4-Dimethylketone-3,4,3,4- Tetrahydro- <i>n</i> -Triazin-1-one) (185)	1955
8-(<i>p</i> -Tolyl)-3-Isopropylisothiazolidine (186)	1955
Isomethylketone-8-(<i>p</i> -Tolyl)-1-one (186)	1955
Alkylation of Dimethylketone-8-(<i>p</i> -Tolyl)-1-one (204) with Methyl Fluoroborate, 1,1,3-Trichloro-2-(3,4,4-Trichloro- 2-oxopropylidene- <i>n</i> -Triazin-1-one) and 1,1,3-Trichloro- 2-(3,4,4-Trichloro-2-oxopropylidene-3,4,3,4-Tetrahydro- quinazolin-1-one) (187)	1955
2-Isopropyl-2,3-Dichloroquinazolin-1-one (188)	1955
Hydrolytic Action of Isopropylketone (207)	1955
Alkylation of Dimethylketone-8-Isopropylketone (111) by 2-Isopropyl-2-(3,4,4-Trichloro-2-oxopropylidene-3,4,3,4-Tetrahydro- quinazolin-1-one) Fluoroborate (112, 113)	1955
2-Isopropyl-2-(3,4,4-Trichloro-2-oxopropylidene-3,4,3,4-Tetrahydro- quinazolin-1-one) (114)	1955

Treatment of 8-Methyl-8-oxaethyl-5-Chloro-3-Methyl-3-Propenylamine (137) with Sodium Hydroxymethylsulfonylchloride	134
Treatment of 8-Methyl-8-oxaethyl-5-Chloro-3-Methyl-3-Propenylamine (137) with Potassium <i>n</i> -Butoxide	134
8-Methyl-8-oxaethyl-5-chloro (137) 	134
8-Methyl-8-oxaethylammoniumchloride (138) 	137
8-Methyl-8-oxaethylammoniumchloride-5-Chloro-3-Methyl-3-Propenylamine (138) 	138
Treatment of 8-Methyl-8-oxaethylammoniumchloride-5-Chloro-3-Methyl-3-Propenylamine (138) with Sodium <i>n</i> -Butoxide	139
8-Hydroxyethylsulfonylchloride (142) 	139
3-Ethylpropyl-2-N-propylamine (143) 	139
8-Methyl-3-isopropyl-3-oxopropionamide (144) 	139
Reduction of 8-Methyl-3-isopropyl-3-oxopropionamide (144) with Sodium borohydride. 8-Methyl-8-oxaethyl-3-oxopropionamide (145) 	139
<i>n</i> -Butylamine (46) 	139
8-Methyl-3-oxo-1,3-diphenyl-2-oxalidiammonium (147) 	139
Thermolysis. Isomerization of 8-Methyl-3-oxo-1,3-diphenyl-2-oxalidiammonium (147) to 8-Methyl-3-oxo-1,3-diphenyl-2-oxalidiammonium (148) 	139
Thermolysis of 8-Methyl-3-oxo-1,3-diphenyl-2-oxalidiammonium (148) to 8-Methyl-3-oxo-1,3-diphenyl-2-oxalidiammonium (149) 	139
Addition of <i>n</i> -butylamine (46) to 8-Methyl-3-oxo-1,3-diphenyl-2-oxalidiammonium (149). 8-Methyl-3-oxo-1,3-diphenyl-2-oxalidiammonium-8'- <i>n</i> -butylamine (150) 	139
Acid Catalyzed Hydrolysis of Phenyl-(2-Phenyl-8-Carboxampho- methyl-2-oxalidiammonium-8'- <i>n</i> -butylamine (150). 2-Phenyl-3- (2-Phenyl-8-Carboxamphoethylammonium-8'- <i>n</i> -butylamine) (151) 	139
Phenylsulfonyl Chloride 	139
1-Propan-2-Phenylsulfonyl bromide (152) 	139
1-Propan-2-Phenyl-8- <i>n</i> -butylamine (153) 	139
Triethyl 8-Phenylglycinate (154) 	139
Independent Synthesis of 2-Phenyl-3-(2-Phenyl-8-Carboxamphoethyl- ammonium-8'- <i>n</i> -butylamine) (155) 	139
EXPERIMENTAL	139

THE SPECIAL	245
GEOGRAPHICAL SECTION	271

LIST OF TABLES

Table	Page
I. 100 MHz Proton Spectrum of Terephthalic Acid <u>10</u>	10
II. ¹³ C Chemical Shifts for <u>10</u> $\rightarrow$ <u>11</u> $\rightarrow$ <u>12</u> $\rightarrow$ <u>13</u> $\rightarrow$ <u>14</u> $\rightarrow$ <u>15</u>	10
III. 100 MHz Chemical Shifts and Coupling Constants for <u>11</u> $\rightarrow$ <u>12</u> $\rightarrow$ <u>13</u> $\rightarrow$ <u>14</u> $\rightarrow$ <u>15</u> $\rightarrow$ <u>16</u>	10
IV. A Partial Tabulation of NMR Spectral Parameters from Observed Data <u>16</u> and <u>17</u> $\rightarrow$ <u>18</u> $\rightarrow$ <u>19</u> $\rightarrow$ <u>20</u>	10
V. 100 MHz Chemical Shift and Coupling Constants for <u>16</u> $\rightarrow$ <u>17</u> $\rightarrow$ <u>18</u> $\rightarrow$ <u>19</u> $\rightarrow$ <u>20</u> $\rightarrow$ <u>21</u>	10
VI. Solvent Effects Concerning the Cyclization of <u>21</u> $\rightarrow$ <u>22</u> $\rightarrow$ <u>23</u> $\rightarrow$ <u>24</u> $\rightarrow$ <u>25</u> $\rightarrow$ <u>26</u> $\rightarrow$ <u>27</u> $\rightarrow$ <u>28</u> $\rightarrow$ <u>29</u>	10

Statement of Dissertation Presented to the Graduate Council
of the University of Florida in Partial Fulfillment of the Requirements
for the Degree of Doctor of Philosophy

RETROSYNTHESIS FROM AND FLUOROPHOSPHATE SALTS
RELATIONSHIP TO NOVEL, RETROSYNTHETIC REACTION

By

George L. Stein

December, 1976

Chairman: James A. Maykop
Major Department: Chemistry

A new route to the theoretically interesting allylic ether substitution reaction was explored. N-*o*-allyloxymethyl-N-allylimine was acetylated with methylfluoroacetate to give the respective ketocarbonate. Chloromethyl-*o*-allyl ether was also in high yield. Acetylation of this salt with sodium bis-(2-allyloxyethyl)phosphate in benzene failed to afford the anticipated chloromethyl ether due to the allylic ether substitution via a bimolecular yield. Instead, elimination to 2,2'-bis(2-allyloxyethyl)-1,3-bis(2,2,4,4-tetraallyl-3-oxocyclopentyl)propane occurred, presumably by a $\pi^4_s + \pi^4_s$ spirocyclic of the 1,2-diphenyl methylene. Experimental evidence for the interaction of a ketocarbonate yield is given. Spectroscopic and mechanistic interpretations are also discussed.

The allylation of ketocarbonates was bearing especially promising N-allyl groups on the ketone carbonyl gave different results. In contrast with methyl fluoroacetate, N-allyl ketocarbonate produced chloromethyl ether with sodium perchlorate-*o*-chloride, depending upon the

sign of the minimum varied on the basis of frequency. 3-Allyl
indolizines gave only 3-indolizidinone fluoromethide salts.
Spectroscopic data and mechanistic interpretations are discussed.


The behavior of indolizines under isomerization is described
in most cases, addition or decylation products were obtained. Mechan-
isms consistent with these transformations are described.

A system theoretically inclined to undergo a rare 3 β cyclization
was investigated. The addition of indolizidinone to ethyl trans-5,8-
dihydroindolizidinocyclopentane was effected in solution with carbon
tetrachloride. The expected 3 β cyclization product was not observed,
instead, phenyl-(8-phenyl-5-oxo-8-oxapentyl)indolizidine-8 β -1-benzylidene
was obtained in excellent yield. These acid anhydrides proved to behave
tetrachloride were degraded in the high conversion to the indolizidinone.
Mechanistic details are discussed.

CHAPTER I
 KINETICS AND MECHANISM OF
 11-4-EPOXYDATION OF BUTYL-6-CYCLOHEXYL
 PLASMO-POLYMER


Introduction

isobutylene derivatives of the general structure 1 and their oxirane head isomers 2 and 3 are relatively obscure compounds. They have only recently attracted the attention of synthetic chemists




because of their analogy to the methylene cyclopropane system, and hence their potential for molecular rearrangement.¹⁻⁴ Indeed, Quast and Maxam² observed the thermal rearrangement of the isopropylidene derivative 4 to the isobutylene cyclopropane 5 as illustrated in Scheme 1.

SCHEME 1



ring action of the methylene aziridine, since these systems failed to undergo exchange with solvent under typical conditions, an alternative addition mechanism of the type depicted in Scheme II was proposed.

SCHEME II




Finally a decade later, Dwyer and Greenwald⁸ reported an unusual rearrangement of aziridine 11. When refluxed in *n*-butanol with an excess of potassium *n*-butoxide, 11 was converted to a product identified as hydroxycyclopropane 12. It was suggested that the rearrangement was initiated by deprotonation of the aziridine 11 to give the methylene aziridine 13 which subsequently isomerized to the cyclic cyclopropane 12 under the experimental conditions. A relatively straightforward mechanism was proposed by the transformation of 11 to the anion 14.^{9,10}

Kricheldorf and Bachter¹¹ attempted to provide a direct synthesis of substituted aziridines by the addition of azirones 15 to the thermal or electrolytic decomposition of electronegatively substituted oxides to alkenes. In most cases, 1,3-dipolar addition to the double linkage of the alkene took place with the formation of 1,3,3-triazolones 16. Thermal decomposition of these triazolones, which in other systems were known to produce aziridines and alkenes,¹² gave only α,β -unsaturated

SCHEME III



Substitution C_{25} . In our system, thermalolysis of vinyl sulfideformate II in refluxing tetramethylsilane I produced in addition to trioxolane III , a compound which had spectral properties consistent with compound IV .¹⁵ The origin of IV was not unequivocally established. Two possible routes




of attack by the nitron across the alkene were suggested: a 1,4) dipolar cycloaddition of carbonyl nitron to the alkene, or a 1,4) addition to give intermediate VI which then rearranged via a 1,5) sigmatropic decomposition under the experimental conditions. In no instance was direct evidence for an allylic nitron addition given.


More recently, Kingston and Moffatt¹⁶ have reinvestigated the addition of vinyl sulfideformate II to substituted alkenes. Here catalyzed isomerization of 5-(p-methoxycarbonyl)pentene was used as a source of carbocationic. Upon treatment of a dilute solution

chloride anion of 1,1-dimethylallene 22 with carboxylic anions,

SCHEME IV




an unstable chloroalkene **22** was obtained in approximately 70% yield. Spectroscopic analysis¹³ of this substance allowed assignment of the 2-isopropylidene-2-methylcyclopropanol **23** structure. Intriguingly,



photolysis at 130° of a dilute solution of **23** in carbon tetrachloride produced the carotenes **24**. This was identical to the carotenoid obtained by Haxelbauer and Staehle¹⁴ for the carotenoid formed from the photolysis of vinyl caradiene **25** to 1,1-dimethylallene **26** (Table 1, entry 1).


The use of isopropylidene as synthetic precursors of allylic caradienes has been the subject of study in the recent literature. Cornhill and Crouley,¹⁵ for example, treated 2-isopropylidene-2-methylcyclopropanol



with a variety of methylene transfer reagents including diazomethane, dimethylketene acetylen and dimethylacetone acetylen. Addition of diazomethane to isocyanide II proceeded sluggishly, and afforded methanone add II in low yield as the only identifiable product. Reaction of II with sulfur phos proceeded readily, but




again unsaturated products were obtained as illustrated in equation (7) and (8). The earlier reports concerning the addition of carbene



to isocyanides showed addition occurring across the carbon-nitrogen bond rather than the carbon-carbon bond as observed by Espada¹² and Campbell.¹³ Jandry and Goshaw¹⁴ reported that diazomethane added

usually to diphenylketene-*O*-(*p*-bromophenyl)idene in yield > 1,1,1-trichloro-2,2,2-trifluoroethane. Indeed, and this,¹⁷ on the other hand, demonstrated that addition of dichloromethane to diphenylketene-*O*-(*p*-chlorophenyl)idene **24** did not give the expected product **25**. Instead, a compound was isolated with elemental composition and spectroscopic properties in agreement with the bisdichloromethane adduct **26**.

SCHEME V




Because of the limited generality of synthetic routes to the allylidene acetalides until then far presented in the literature, a new pathway to this interesting system was explored. The problem was approached by taking into consideration the chemistry of functionalized acetalides, of which allylidene acetalides are members, and their ability to undergo thermally allowed ring opening to immediately yield furans. Indeed, Myers and Rhee¹⁸ created a variety of aldehydes and ketones with ortho functional groups and obtained respective furan derivatives in high yields. In one case, diphenylketene with a carboxyphenyl group has produced acetalide **27** in quantitative yield,


possibly via the anion 14.

This research was directed toward the generation of a boronium ylide of the general structure 15 via alkyloxy boroninium salts 16.

SCHEME VI



with the intent of directing cyclization to the corresponding allylic enol ether 13. This approach proved particularly promising not only



because of the ability of anionides 14, but also due to the striking resemblance to the mechanism (Scheme II) proposed by Ingold and Mann⁸ for the cyclization of *N*- β -hydroxyethylchloracetate to methylene sulfolene.

Although allyloxy boroninium salts are not known, solutions of chloroborate salts from the treatment of chloroboranes with allylic chlorohydrates have been proposed.¹⁰ These solutions proved to be

extremely sensitive to electrophilic substitution on conjugated aromatic systems,¹⁰ (vi) copolyoligomers with alcohols^{10,11} and aldehydes,¹² and (v) general unacceptably hygroscopic and insoluble limited attack by nucleophilic reagents.¹³ The high reactivity, or conversely the low stability of these ketosidation fluorosiloxane walls is explained in part by the size of hole created by the substituents. It is known, for example, that shorter alkyl ketosidation are comparatively stable under acid or base melting agents, long alkyl ketosidation are thermally unstable with respect to polymerization, and tend to hydrolyze readily in acidic media.¹⁴ All of the ketosidation fluorosiloxane walls presently known are ketosidated with groups no longer than ethyl. As a result, the small volume occupied by these substituents allows virtually no resistance to nucleophilic approaching the highly conjugated sp²-hybridized central carbon of the ketosidation wall. These walls for various reasons show no tendency to polymerize, but otherwise are perfect electrophiles.

The selection of a suitable ketosidation wall prior to the synthesis was, therefore, subject to certain structural restrictions. First, the substituents had to be large enough to stabilize the derivative, and subsequently the conversion to a three membered ring.¹⁵ Subsequent efforts of this nature were implicated in the remarkable stability of di-*tert*-butylsiloxane $(\text{C}_4)_2$ ^{16,17} which most effectively becomes




4.4

which dissolves rapidly and readily accepts nucleophilic reagents.¹⁸ Before **4.4** was easily distilled as a stable monomer and showed a marked

resonance bond attack by nucleophiles.¹⁶ Recently, these substituents have been shown to be inert to the strongly basic conditions¹⁷ used in alkene isomerization to the corresponding ylid 12. Ideally, the substituents, X' , should be reasonably bulky and nonconjugative in order to discourage attack at the internal carbon under otherwise favorable conditions or chemical conversion of 11 to 12. Scheme VII illustrates the two proposed pathways to be investigated in this work. The direct OR involving the direct alkylation of ketone-alkene 11. Alkylation in general can be accomplished from secondary amide substrates by a variety of straightforward methods.¹⁸⁻²²

SCHEME VII



Elaboration of these methods and their scope will be discussed in Chapter II. The direct alkylation of ketone-alkenes has neither been achieved nor attempted. Furthermore, their relative stability under such strong alkylating conditions as methyl trifluoromethane has yet to be tested.

The treatment of cyclohexenones 14 OR with alkyl fluoroborates (Table, page 4) and their behavior under deprotonation conditions will be discussed in a later chapter.

Results

The sterically protected di-*n*-butylphosphine-B-triethylamine **18** was synthesized via the procedure **11** through **28** as depicted in Scheme VIII. The sequence **11** through **28** was carried out according to Jones' literature procedure^{10,14} while the sequence **11** was accomplished as a partial objective in this research. The conversion of acid chloride **14** to the triethylamide **12** took place smoothly (85%) in liquid nitrogen despite the reported¹⁴ rigorous conditions needed in benzene **18** in the primary amide with sodium azide in liquid ammonia. Amide **11** was treated with one equivalent of phosphorus pentachloride under the mild conditions of refluxing benzene. The isobutyl chloride **11** was not isolated. Instead, it was dehydrochlorinated directly with triethylamine to a mixture (90% P-DE) which contained isobutene **11** and a substance believed to be di-*n*-butylphosphineoxide.¹⁴ Reactions **18** exhibited a gross loss in yield (31,48%) (calcd for $C_{12}H_{24}P_2O_2$ 48,48%) in the mass spectrum. The appropriate signals and multiplicity were obtained in the mass spectrum as well as a strong and very characteristic infrared band at 1198 cm^{-1} associated with the heteroatomic functionality, (note^{14,17}). Attempts to isolate an analytical sample of **18** completely free of water either by conventional distillation techniques or by column chromatography resulted in only slight purification. Nevertheless, the alkylation was performed by cycling a tenfold excess of methyl fluorosulfonate¹⁸ into a stirred ethanol solution containing isobutene **11**. Alkylation occurred rapidly and selectively at the isobutene nitrogen, resulting in an almost immediate precipitation of the isobutylphos fluorosulfonate salt **19** as a white fluorosulfonate solid. This material was determined by spectroscopic

analysis to be completely void of nitrite under dehydrated nitrite by-products. Isocyanidation only 10 proved to be remarkably stable (up 150-160° with decomposition) considering its molecular environment and the known chemistry of other heterocyclanes.^{12,13,14} It has a high degree of solubility in polar solvents such as chloroform, ethanol, or water, and can be recrystallized from methylbenzylchloride/ether. Surprisingly, it is resistant under neutral hydrolytic conditions to the point where it can be recovered unchanged after sitting in aqueous solution at room temperature for two hours or more! The infrared spectrum of 10 shows a band of medium intensity at 1600 cm^{-1} for the isocyanidic functionality. This band seems to be at somewhat lower frequency than expected for a isocyanide with a positively charged heterocycle. In NMR data, the methyls, also appreciably influenced by this is higher frequency upon protonation or alkylation.¹⁵ Also, the higher frequency of isopropenyl¹⁶ (10, 1675 cm^{-1}) and ketone¹⁷ (10, 1708 cm^{-1}) compared with that of 10 tend to make the band at 1600 cm^{-1} appear relatively low. Present in the nmr spectrum is a low field olefinic signal at δ 1.59 and a deshielded methyl signal at δ 1.95, as well as the expected olefin protons at δ 1.98 (olefinic) and δ 4.11 (quartet). As further structural proof, 10 was hydrolyzed in aqueous base to the tertiary amide 11 according to equation (11):



The structure of this anion was established by high resolution mass spectrometry with a parent ion at m/z 113.059 (calcd for $C_{13}H_{20}SiO^-$ 113.059), and by ^{13}C NMR and mass chromatograms.

The degradation of **10** was performed in toluene using sodium hexafluoroantimonate as a sterically hindered, non-nucleophilic strong base.¹⁴ Stirring a slurry of **10** in toluene with sodium base for 24 hours produced the piperidine ester **12** in 12% yield rather than the intended acrylate **11**. The linear structure of **12** was confirmed by the high resolution mass spectrum with a parent ion at m/z 186.103 (calcd for $C_{21}H_{30}N_2O_2$ 186.103).

SCHEME IX



The partial non-equivalent anisotropic acetylation protons (H_A , H_B) of piperidine **12** appeared as an AB pattern in the run with coupling constants and chemical shifts data shown in Table I.

Table 1
180 Mc. Proton Spectrum of
Diphenyl Ether 41



41

Proton	δ	Multiplicity	
H_1	6.87	singlet	$J_{1,2} = 0.7$ Hz
H_2	6.80	singlet	
H_3	6.75	singlet	
H_4	6.80	doublet	$J_{4,5} = 10$ Hz
H_5	6.75	doublet	$J_{5,4} = 10$ Hz
H_6	6.62	doublet of quartets	$J_{6,7} = 11$ Hz, $J_{6,8} = 7$ Hz
H_7	6.72	doublet of quartets	$J_{7,6} = 11$ Hz, $J_{7,8} = 7$ Hz

The aromatic methylene protons (H_6 , H_7) displayed an interesting quartet of quartets pattern which collapsed to a single AB system upon spin decoupling of the methyl protons (H_8). The partial non-equivalence of these protons (H_6 , H_7) can be attributed to restricted rotation of the *o*-methyl groups of 41.⁴¹ Inspection of molecular models clearly shows a considerable degree of steric interaction between the *o*-methyl substituents and the neighboring methyl group. Also, the vinyl


intensity according to 1:1 in (vs. 0-511 ppm) identified relative to the vinyl signal, or 1:1. It could conceivably be the result of simple anisotropy which is known to cause paramagnetic shifts of this magnitude.¹²

When compounds appearing in the degradation product analysis showed infrared absorption bands of rather weak intensity at $\mu\mu$, 1000 and 1646 cm^{-1} , these were interpreted as arising from decarboxation and hydrolysis of the ketocyclization salt II_2 during the prolonged reaction times. Decarboxation has been noted by Heyrup and Smith¹³ to be an anapoly side reaction, but in general, could not be entirely eliminated. Partial hydrolysis of II_2 was expected to result due to traces amounts of moisture in the solvent, and/or hydrolytic contaminants present in the allylamine base.

Attempts to convert ring closure by performing the isomerization in refluxing benzene again produced epimerate II_2 and in somewhat higher yield (50%). Isomerization under the homogeneous conditions of triethylphenylammonium (20%) resulted in a drastic decrease in diastereomer (III), but only at the expense of hydrolysis to the acids II_1 (III). Further attempts to effect cyclization were abandoned, and attention was focused on the identity of the supposed "1,3-dipole" structure of II_2 .

Chemoselective additions to 1,3-dipolarophiles have been of profound importance in establishing the identity of azabicyclic ylides during the course of azabicyclic isomerizations.^{4(b-d)} These trapping experiments are now recognized as constituting evidence for the intermediacy of other 1,3 dipoles as well. Unfortunately, reports of in situ generated 1,3-dipoles by conventional trapping agents was

subject to major experimental problems. Most desirable diphenyls¹² would readily sublimate. The necessity of the strongly acidic conditions required for dipole formation, however, was found to be completely inert to the allylolside bond under the decomposition conditions. Treatment of **12** with sodium bis(trimethylsilyl)fluoride in the presence of a tenfold excess of methanol **13** during the 1.5 minute **13**-¹⁹ treatment.



purification by short path distillation resulted in some sample decomposition with no substantial improvement in product quality. Spectrometric analysis of sides **13** before distillation revealed a pattern in the IR (IR) (solid film $C_{20}H_{24}O_2$) IR (IR) Complete analysis of the same spectrum suggested the presence of at least one other component; the identity and relative percentage of this by-product(s), however, was not determined ($C_{20}H_{24}O_2$). NMR analysis showed the expected nonaromatic α -methyl signals at δ 1.35 and 1.45, a methyl triplet at δ 1.80 as well as unreacted methylene and methoxy signals in the range δ 1.0-2.0. ¹³C NMR (Table II) proved to be helpful in establishing the placement of the olefinic linkage carrying the α -methyl groups. These appeared as singlets at δ 141.4 and δ 134.8 downfield from tetramethylsilane and are in agreement with typical shifts for sp^2 hybridized carbons. The assignments made in Table II were based on the multiplicities observed from single frequency off-resonance decoupling, these together with typical chemical shift values for vinyl methoxy and pyridine derivatives.¹³ None of these assignments are

Table 11

^{13}C chemical shift data for **55**




55

δ^{a}	Assignment	Assignment
16.5	quartet	1
16.7	triplet	8
11.1	quartet	16
11.6	triplet	2
10.6	quartet	15
10.1	singlet	14
10.0	singlet	13
10.7	doublet	3
11.1	doublet	9
11.3	triplet	10
16.3	triplet	7
11.1	doublet	4
14.6	doublet	6
17.4	triplet	5
11.8	singlet	11
110.8	singlet	12

^aValues in table are reported downfield from TMS with deuteriochloroform as a solvent.

be in error due to small differences in chemical shifts, but, for the most part, the spectrum fits well for the proposed structure 41. The infrared spectrum exhibited a weak band at $990-1020\text{ cm}^{-1}$ characteristic of the trans double bond.²⁸

Interestingly, when 41 was repeatedly distilled during an attempted purification 41 failed chromatography. After several elutions through an alumina column with chloroform or methylene chloride a substance was obtained whose spectral properties and elemental composition were in agreement with 41. This same product was isolated from a chromatographically acid treatment of a distillate containing 41. Presumably, the small amounts of hydrogen chloride known to be present in chlorinated hydrocarbons were responsible for the polymerization during chromatography.



The new spectrum of this derivative was much more informative than that of 40. The positively charged nitrogen greatly deshielded the adjacent pairs of methylene protons while separating the H_{ax} and H_{eq} gem protons by anisotropically δ/π use. Table III provides chemical shift values and a partial list of coupling constants for 41. The gem configuration of which 41 was established by the magnitude of the $\text{H}_{\text{ax}}-\text{H}_{\text{eq}}$ gem $\text{H}-\text{H}$ proton coupling constant (3.5 Hz), and the angle literature presented for the preferred gem addition of 1,2-dipoles to carbonyles.^{24,25} In general, the gem-gem coupling constant by itself

Table 18


Partial Tabulation of Mass Spectral
Fragmentations Obtained from 15 and 17

1517

m/e	Relative Intensity (%)	
	<u>15</u>	<u>17</u>
100 (M^+)	1.4	1.8
104	3.4	3.3
120	11.3	10.6
132	18.6	19.6
148	17.3	16.7
166	100.0	100.0

these trends which by a variety of means including methyl addition (oxidation and/or deprotonation), carbon-hydrogen bond cleavage (oxidation), carbon-hydrogen bond cleavage (oxidation), and aqueous carbon-hydrogen bond cleavage (oxidation and/or hydrolysis). The unusual prevalence of methyl 15 in samples 15 and the marked prevalence of 17 toward deprotonation suggest further common structural analysis of 15, 16, and 17 molecules which show that extensive crosslinking between the α -hydroxyl groups and the carbon-carbon double bonds in

SCHEME 8




unavoidable (51, Figure 2). Substituted norbornenes, though, as experienced in previous work reaction of the carbon bearing the *t*-butyl group causes a thermodynamically more stable conformation (52, Figure 2) as illustrated in Scheme 8. The resistance of 51 to

SCHEME 9




deprotonation should not be obvious; but proton abstraction to convert the carbon bearing the *t*-butyl group must result into a sterically demanding conformation so that the proton is in forced closer a particular arrangement with the π -orbital of the leaving bond. This




unsaturated feature results in its reluctance to respond under deprotection conditions. The strong toxicity of 1,4-bis(benzyloxy)-aminopyrrolidines,¹¹ for example, has been attributed in part to relief of strain accompanied by protonation.

5-Methyltetrahydrothiazolines 18 was used as a 1,2-dipoleophile with hopes of simplifying the previous reaction sequence, and possibly producing a crystalline tetraolamine adduct 19. The acetal species 17



was again prepared as an oily mixture and could not be isolated. Rather, the hydrochloride salt 19 and its isomer 19' were obtained and characterized as solids. ¹³C analysis clearly shows the expected presence



of isomers 19 and 19' as a matter of time pulse, particularly in the aromatic region δ 119-120. The signals associated with the minor isomer, presumably 19', were based entirely on the selective intensity of each pair of lines. The lower intensity lines being assigned to the minor isomer. The 120 MHz proton and spectrum of 19 confirmed the *syn* configuration from coupling constant parameters. These values were,

as expected, very similar to those observed for $\underline{\text{X}}_1^-$. Because of the fused aromatic ring, a more complex analysis of coupling needs to be made. The results of this analysis are listed in Table V.

Discussion

The experimental observations of dimerization and nonspecific addition to uracines can best be explained in terms of the ionization of $\underline{\text{X}}_1^-$. Dimerization of 4,5-dipyridine is not new, and several reports




of them are found in the literature.¹²⁻¹⁵ The mechanistic pathways accounting for dimerization can be developed: concerted 2d⁰ sp² sp² addition resulting from orthogonal approach to two $\underline{\text{X}}_1$ and a concerted two step addition. These are illustrated in mechanisms a and b respectively in Scheme III. Resonance forms of each species show an appreciable difference in the amount of the delocalization experienced in their transition states. Even so, mechanism b seems

SCHEME III

Mechanism a




Table 7
 900 MHz (CDCl₃) and Coupling Constants
 Data for **20**



20


Proton	δ	J	Calcd (Hz)	Found (Hz)
$H_{1,2}$	7.80	$J_{1,2}, J_{1,3}^{+2}, J_{2,3}$	6.4 ^{40,42}	4.9 ⁴²
H_{3a}	7.80	$J_{3a,3b}$	6.8 ⁴²	6.0
		$J_{3a,3c,d,e}$	7.5 ⁴²	10.0
		$J_{3a,3c,d,e}^{+2}, J_{3a,3c,d,e}^{+2}, J_{3a,3d}$	15.0 (1.0) ^{40,42}	11.8 ⁴²
		$+J_{3a,3b}^{+2}, J_{3a,3d}$		
H_{3b}	7.90	$J_{3b,3a}$	6.8 ⁴²	6.0
		$J_{3b,3a}^{+2}, J_{3b,3c,d,e}^{+2}, J_{3b,3d}^{+2}, J_{3b,3e}$	12.2 (11.5) ^{40,42}	14.0 ⁴²
H_4	7.94	$J_{4,3a}^{+2}, J_{4,3b}$	12.1 (11.5) ^{40,42}	3.0 ⁴²
H_5	7.94	$J_{5,3a}^{+2}, J_{5,3b}$	12.1 (11.5) ^{40,42}	3.0 ⁴²
$H_{3c,d,e}$	8.0-8.07-8.01			
$H_{3c,d,e}$	8.02	$J_{3c,d,e,3a}$	6.1 ⁴²	10.0
		$J_{3c,d,e,3b}$	11-13 ⁴²	11.0

Reaction 3:



selectivity. Considering the relative concentrations of the diols 21, 22, the bromination with 23, a competing attack by the diols on 23 should become a competing process if not the major one. No products were detected which could lead support to this mechanism. Reaction 4, therefore is favored.

Transition-State stabilization fluorescently cyclizable in solution with exceptional ease is what has been described as a $\pi^*_A + \pi^*_D$ concerted process.²⁵ Like before,²⁶ the ability to accomplish this is due to strong bonding interactions between the electrophilic halogen cation and the π system of the olefin. The absence of steric hindrance and transition state strain inherent to typical olefinic systems are contributing factors, too.²⁷ It is logical to assume that this case




electrophilic bonding interaction is present in solution is the proper geometric orientation of piile 21 prior to bonding. The electrophilic

interaction in this case would be with the nitrolic terminus of the heteroleptic pldd II. This overall concerted process, then, could be classified as $\sigma^2_{\text{N}}\pi^2_{\text{O}}\sigma^2_{\text{C}}$ cycloaddition.

The failure to observe electrophilic ring closure was perplexing on first scrutiny, since the requisite dipole almost certainly had been generated. A rationale for this apparent anomalous behavior is offered in Scheme III. For cyclization to occur, the dipole in either of the two valence bond resonance structures must successfully pass through a transition state whose electronic environment can formally be described as II or II'. In either case, a high energy carbonium ion center¹⁸ must be maintained throughout the cyclization until sufficient energy can be added with the nitrolic carrying the


(SCHEME III)



negative charge. Although direct analogy from carbocation intermediates was not found in the literature, it is known that substituents which can stabilize the incipient anion which will facilitate the rapid reversible ring opening process. Substrates lacking such substituents tend to undergo carbon-carbon or carbon-hydrogen bond scissions and polymerize when subjected to thermal ring opening conditions.²²

It would be of interest to design such in this case to consider the synthesis of heterocyclic acids bearing potential glycid stabilizing groups, particularly those incorporated as nitrogen substituents. An example system, for example, could serve a useful function; i.e. QD to directly assist in the ring closure via conjugative stabilization of the glycid, and QD to enhance the acidity of the protons adjacent to the incision nitrogen.

SCHEME XIV




CHAPTER III
KINETICALLY FAVORABLE FREE-
RADICALIZATION OF ACETYLENE BY
METAL FLUOROBORATES

Introduction

In Chapter I, it was shown that a sterically protected tertiary ketoxalate experiences facile and rapid allylation with vinyl fluorosulfonate to produce a stable, reprecipitable ketoxalate-fluoro-sulfonate salt in high yield. Different results were obtained, though, when the bulk of the ketoxalate substituent was reduced. In these cases, diene and triene were formed. Barker and Bennett⁶²⁻⁶³ have recently reported the thermal chemistry of various ketoxalates. They suggested that specificity for diene and triene formation was found upon allyl, β -aryl ketoxalates. For example, ketoxalates which thermally eliminate to any appreciable extent are the fully aromatic substituted ketoxalates. In contrast, 1-ketoxalates appear to arise from diarylketone- β -allylketone (ketone:O%₂). Species with β -aryl functionality, unsubstituted in the α position, yield quinodienes (diene) as the major product. No mechanistic conclusions or speculations were given by Barker in his publications.

The objective of this chapter, then, is to describe the allylation behavior of several ketoxalates under strong allylation conditions. Chemically, this work provided the evidence obtained in Chapter I. However, it was the indicative chemistry and mechanistic implications to be discussed in this chapter that led to the eventual formation of the reprecipitated ketoxalate yield 24. An attempt was made to

SCHEME XV



recommendations the specificity or selectivity of ether and ester formation in terms of substituent effects, thermodynamics, and mechanism. These questions will be correlated with those reported by Barker, and possible mechanistic pathways will be discussed. Also, the use of methionemimides

will be evaluated as sources of bromination sites, and, thereafter, processes to allylic bromination through π - π -bipolar intermediates.


EXPERIMENT

The bromolactams used in this work were prepared from their secondary amides according to a variant of the published procedures depicted in Scheme IV.¹⁸⁻²² The particular bromination substitution patterns needed for individual experimental design resulted in systematic synthesis, when a wide variety of carboxylic acids and primary amides for the respective amides were commercially available.

SCHEME IV



Alkylation of a dilute ethereal solution of *di*-*tert*-butyl-*trans*-2,3-phenylidene **12** with methyl diazoniumfluoroborate proceeded slowly. In



immediate color development, generation of heat, or substantial noticeable changes in the solution were observed until the first appearance of a deep colored precipitate usually 10 minutes after the introduction of the methylating agent.

The nmr spectrum of this precipitated material showed a decoupled methyl resonance at δ 1.18 in addition to a pattern of aromatic signals. Integration of the methyl and aromatic signals revealed a dimeric structure, and suggested a 2:1 adduct between hexafluoroborate **12** and methyl diazoniumfluoroborate respectively. The presence of the diazoniumfluoroborate counterion (MeN_2F_6^-) was established by the appearance of delayed absorption maxima at 1215 cm^{-1} and 1045 cm^{-1} . These bands have been associated with the asymmetric and symmetric sulfur-sulfur vibrational modes characteristic of sulfinic acids.¹² The nmr spectrum exhibited



a paralet period (as at c/c 112 ϕ^2 - HNO_2/P) with no major frequency changes other than the observed loss of 12 mHz units recorded in the cleavage of the *N*-methyl group. Spectral data and electrostatic calculations supported the quinonoid structure 41, thus this was not sufficient basis for structural assignment, attempts were made at chemical degradation. Treatment of 41 with a 20% sodium carbonate solution in dioxane produced in 80% quantitative yield a brilliant fluorescent yellow compound. The high resolution mass spectrum of this material showed a parent ion at m/e 166.2181 (calcd for $\text{C}_{14}\text{H}_{10}\text{N}_2\text{O}_2$: 166.2164), and a fragmentation pattern identical to fluoranthene with 42. The net spectrum confirmed an upfield shift of the *N*-methyl signal from δ 3.56 to δ 3.48 as expected from migration of the positive charge from the quinonoid ring 41. This data suggested simple loss of the dioxane HNO_2/P leaving the base unchanged, and led to the structural assignment 43.




Treatment of 41 with sodium borohydride proceeded rapidly and quantitatively to 44 which exhibited an IR spectrum centered at δ 4.92. The *N*-methyl signal at 3.48 was found to be shifted approximately 0.3 ppm upfield from the position of the *N*-methyl signal in 41. NMR responses suggest that this shift is too large to arise from solely an increase in the anisotropic deshielding. The origin of this difference is undoubtedly the partial removal of deshielding shielding by the

partial aromatic ring. Identification of molecular orbitals clearly shows that a greater degree of base to base interaction between the methyl group and the aromatic ring can be achieved in II (relative to II'). This argument is further substantiated by the spectral properties of a chelate quinonoid having no aromatic functionality in the 8-position. In this system (vide infra) the 8-methyl resonance appears at δ 3.45.



The reduced quinonoid derivative II proved to be an ideal candidate for an unequivocal structure identification via acid hydrolysis. Vigorous hydrolysis of II in refluxing aqueous acid produced the *n*-octane ketone II' in a 90% yield in 70% yield, which, and trace amounts of dithienylacetophenone. Also, compound II was isolated as a pale yellow crystalline solid in approximately 44% yield. The presence of

SCHEME XVII



this material is presumably the result of uncomplexed intramolecular substitution by diphenylcarbamate on the reaction between $\underline{R}_1$ followed by dehydration. The structure $\underline{R}_2$ was established from the spectral properties. High resolution mass spectrometry of $\underline{R}_2$ showed a parent ion at m/e 475.0312 (calcd for $C_{22}H_{23}NO$: 475.1644). A base peak corresponding to the loss of a benzyloxy fragment appeared at m/e 231, a low frequency carbonyl stretching vibration attributed to intramolecular hydrogen bonding appeared at 1695 cm^{-1} in the infrared.^{20,21} This value is slightly higher than the carbonyl band for $\underline{R}_1$ which is shifted 12 cm^{-1} lower to 1683 cm^{-1} . The nmr spectra of both $\underline{R}_1$ and $\underline{R}_2$ exhibit doublets (1H) at δ_{ppm} 7.1-8 which collapse to singlets (2H) when treated with H_2O , and broad singlets (1H) at δ_{ppm} 7.5-8 which completely disappear on similar treatment with H_2O . The extremely low field H-H resonances (δ_{ppm} 7.5-8.5) are considerably lower than most reported aromatic aromatic ketones,²² and are reflections of the strong intramolecular hydrogen bonding with the carbonyl oxygen.²³ A rationale for the abnormally low value of the H-H protons is not immediately obvious. Steric crowding between the benzyloxy and the other ketone phenyl rings may cause the carbonyl group to "bow in" toward the other substituent. This would allow for stronger hydrogen bonding and possibly account for the observed low field H-H resonances. This interpretation also explains the observed H-H coupling with the benzyloxy protons which is the reverse




responsible for the C_{β} -methyl shift. In all likelihood, this same phenomenon is operating in 12, but, unfortunately, overlapping aromatic signals make accurate assignments very difficult.



Isocyanides carrying aromatic functionality only as nitrogen were examined. These were expected to undergo cyclization to quinazoline derivatives as did the tertiary isocyanides 13 and 14. Support for this anticipated behavior appeared in the literature ⁴¹ while this work was in progress. The principal reason for examining these isocyanides was to aid in the eventual accurate elimination of quinazoline derivatives 15 from 16 which prior to this work were not firmly established. It was reasoned that a great portion of the analytical spectroscopic data of the tertiary isocyanides appears was being obscured by the presence of "noise" aromatic signals. Further clarification could be obtained by replacing the phenyl substituents on the isocyanide groups by alkyl groups.




Thus, dimethylcarbamoyl-8-phosphorin 17 was prepared as a greenish yellow oil from the nitrile precursor 18. Treatment of a dilute solution of this isocyanide with methyl fluorosulfonate produced after 24 hours a clear solution over a small amount of brown oil. The clear



aldehyde was separated from the oil and evaporated to a colorless crystalline solid (mp 165-166°). The mass spectrum of this compound showed major peaks at m/e 433 (indicative of a trimer) and 145, and a strong infrared absorption maximum at 1615 cm^{-1} - a higher ester frequency associated with 1,45 for the vinyl position. This same trimer could be obtained in 85% yield by treating a dilute ethereal solution of 2-mercaptohexano-2-phenylthioether 15 with a catalytic amount of diethyl-sulfuric acid. Blegel, Smith, and Heston⁴⁵ reported a benzothiole trimer (mp 155-156°) isolated from the photolysis of 15 and various in a β -substitution synthesis. The trimer, identical in every respect to the one obtained in this work, could not be produced by unassisted photolysis or thermolysis of the benzothiole monomer. The compound was believed to be 16 or 17, but the authors claimed that the spectroscopic data alone was compatible with either structure. The system

1617

was not investigated further. In a later paper, Ingber and Thomas²¹ assigned structure 122 to the isomer. Their arguments were based heavily on spectroscopic comparisons of "steroidal analogues" systems and those of a steroid 19 derived from the chemistry of dihydroflavone-8-epoxideflavone 17. Nevertheless, the isomer obtained in this work under alkylation conditions was assigned structure 121. It was found that if the steroidal alkylation product mixture containing 122 was exposed to atmospheric moisture, the yellow crystalline trioxolane diacetonide 123 was produced in 81% yield. This same product could be obtained as a diacetonide 123 by treating a carbon tetrachloride solution of isomer 122 with diacetonide acid. The mass




spectrum of 123 showed a gamma proton loss at ν_{max} 318 ($\nu^{\text{H}}-\text{OCH}_2\text{C}_2\text{H}_5$) and a frequency peak identified to that of 121. A weak absorption band at 1598 cm^{-1} characteristic of most acetones²² appeared in the infrared. The nmr spectrum in trifluoroacetic acid displayed a high field doublet at δ 6.74 [2 ν Hr, 8H], two olefin proton singlets at δ 1.40 and δ 1.26, a multiplet at δ 1.18 [2 ν Hr, 10H], and aromatic protons in the region δ 7.1-7.9. A molecular model of 123 shows that the conformation needed to provide the upfield shift of the isopropyl and isopropylidene methyl protons is easily achieved. The aromatic rings are virtually forced into a disposition orthogonal to the trioxolane ring as a result of steric repulsions with the nearby methyl groups




this shielding factor also applies in the high field resonance of the isopropylidene methyl protons (1.44 δ found in 12). And, it is most likely partially responsible for the anomalous structural assignment made by Fisher and Bennett (1969, 1970).

Fluorobenzene with 12 was easily removed with sodium benzophenone as triazine 13. Alternatively, 12 could be converted back to the parent triazine 11 simply by stirring under an aqueous reflux.



The occurrence of the isopropyl and isobutyl group protonation and the reversible behavior toward protonation and deprotonation leaves little doubt about the structure of 12. However, the simultaneous absence of a quaternization and/or alkylated products in this system was difficult to rationalize. Clearly, the difference in product composition (12) those observed for the nitroalkylation 11 and 13 could only be the result of substituent effects on the kinetic behavior of the intermediates, since this was the only modification made to the substrates. It was not clear,

though, whether the effects governing the cyclization process (steric, π - π interactions) were steric or electronic in nature. In order to clarify this point, chlorophenyl-*m*-*p*-cyclophane 12b with increased bulk in the form of methyl substituents on the ketone functions was treated as an ethereal solution with methyl fluorosulfonate. After approximately



12 hours, the solution was evaporated to dryness, and the poorly soluble residue with ethanol. Residue gave a colorless crystalline solid separated from the mixture. The mass spectrum of this material showed a parent ion at *m/e* 344.0548 (calcd for $\text{C}_{20}\text{H}_{24}\text{O}_2$, 344.1600), and a base peak at *m/e* 187. An aromatic stretching vibration appeared in the infrared spectrum as a medium intensity band at 1448 cm^{-1} . These along with the appropriate signals in the nmr spectrum established the structure of this material as the isomer 12c (20). The ethereal filtrate from which the isomer 12c was crystallized was treated with excess sodium hydride. The quinonoid derivative 12d was obtained from the product almost as quantitative as in the 90% overall yield. The mass spectrum revealed a parent ion at *m/e* 358.0811 (calcd for $\text{C}_{20}\text{H}_{22}\text{O}_2$, 358.1600), and a base peak at *m/e* 338, presumably resulting from the loss of the 2-propyl fragment from the 2-position of 12d. Included in its nmr spectrum were doublet methyl resonances at δ 2.71, 2.47, and 2.31, and a doublet at δ 4.76 (int H_a , 1H, H_b , 1H). In the aromatic region, an ortho (from the 3-methyl group(s)) doublet pattern appeared at

1.4, 32 (benzyl, λ_{max} 268 nm) as well as a low rate of racemization (1.08% epimerization, λ_{max} 268 nm, 18, 55). The relative ratio of oxidation to

SCHEME XIX




epimerization in this system (12.8) suggests that stereoselection is just becoming the kinetically preferred pathway.

From the product distribution observed in the 8-oxyl ketone/ester system investigated thus far, certain trends are clearly evident: (I) systems carrying substituents with bulk appreciably larger than vinyl (e.g., phenyl, *p*-tolyl) favor epimerization preferentially; (II) smaller groups such as methyl tend to direct the cyclization to oxirane derivatives; and (III) groups of intermediate size can be expected to

product mixtures containing quinolizines and mixtures of varying compositions.

Aluminum was found to be a *trans*-alkyl ketonolizine. Here, the possibility of cyclization through a nitrogen substituent could be easily illustrated. In addition, the nitrogen should become somewhat more basic than the *trans*-alkyl derivatives, and, consequently, improve the chances of alkylation and isolation of a monomeric ketonolizine salt.

Hexamethylbenzo-*trans*-quinolizine 111 was prepared by the trihydro-alkylation of the *trans*-alkyl chloride derived from 110. Alkylation of a



chloride solution of 110 with a ten-fold excess of methyl aluminum chloride produced a mixture of 112 and 113. *trans*-Hexamethylbenzo-quinolizine salts 112 and 113 in approximately 50% yield. The




relative ratio of the two isomers was determined by integration of their magnetically nonequivalent *trans*-methyl signals which were found at δ 1.07 (112, 3H, 80%) and δ 1.17 (113, 3H, 80%). The *trans* spectrum was similar in other respects, and, therefore, remains further

interpretation. The 8-methyl resonance at δ 1.12 appears as an unusually highfield resonance to positively charged substituents suggesting 8-methyl anisotropy.^{15,47} This signal is even higher than the 8-methyl resonances observed by Barker and Kucenka⁴⁸ for the diphenylketone-8-methylketone dimer II. The magnitude and direction of this shift is



explainable only in terms of diamagnetic shielding from a nearby aromatic ring, since steric repulsion effects which would consistently be positive would only be paramagnetic shifts. Molecular models of the III isomer III show the phenyl rings in the 8-position properly disposed to offer a diamagnetic shielding environment. This same effect is experienced by the isomeric methyl ketone in the III.



Signal 112 and is observed typically in the upfield area, i.e. 0.800. Apparently, the isopropyl group on the methylene ring adjacent to quaternary C-1 is not affected by the steric strain and shows an unperturbed doublet at δ 1.33.⁴⁸ The high integration of this doublet results from the fortuitous superposition of the olefinic isopropyl resonance from the ggs isomer 111. The effect, then, is a 20% increase over the intensity of a *cis* proton isopropyl pattern. And, as a consequence, the high field isopropyl signal at δ 0.80 of the ggs isomer 112 is only 60% as intense as a full *cis* proton isopropyl doublet 117, Figure 30.





Figure 1: NMR spectrum of 112 and 111,
Isopropyl Splicing Polymers

It is well-known (16) that 111 and 112 were qualitatively reduced with sodium borohydride to a colorless crystalline solid which was identified as the neutral 1-aminocyclohexane 113. Its mass spectrum showed a parent ion at m/e 116.1012 (calcd for $C_6H_{13}N$; 116.1010).



and a lone peak at m/e 101 corresponding to a tetrahydropyridine fragmentation, $(M-15)^+$. The nmr spectrum consisted a singlet at δ 5.42 which was assigned to the methylene ring protons. The 3-methyl and 6-isopropyl methyl signals, however, again were obscured by diamagnetic shielding. The 3-methyl resonance appeared at δ 1.37 while a pair of 6-isopropyl doublets (78 and 82) were found at δ 8.15 and δ 1.15 respectively. Inspection of molecular models clearly shows the ease with which the methyl and isopropyl groups of the cyclohexyl structure can twist about the face of a proximal phenyl group in the 1-position. This accounts for their upfield shifts. Also, a shielding environment for the methinyl 6-isopropyl protons is available provided rotation about the carbon nitrogen bond is completely unhindered. Rotated models, though, indicate considerable steric interactions between the adjacent tetrahydropyridine phenyl ring and the isopropyl methyl group occurs when full rotation is attempted. It is suggested that hindered rotation results in a preferred conformation - as found in the case here - such that one methyl residue is the shielding zone of the ring.

5-phenyl substitution while the other does not. The net effect would permit those of the four isopropyl methyls to experience shielding from



neighboring aromatic systems while the fourth would be relatively unprotected. This analysis is in agreement with the 2:1 integration ratio of the two sets of isopropyl signals. Further evidence for the magnetic inequivalence of the nonidentical isopropyl methyls is found in the NMR spectrum of 11a taken in pyridine (cf. figure 4). Here, three sets of doublets appear





Figure 4. NMR Spectrum of 11a
Isopropyl Splitting 7:4:3:4:6

in the ratios 1:1:2. The two smaller sets previously assigned to the noncyclic and acyclic isomers of the cyclohexene:

Thermolysis of a dry sample of 115 at 350° resulted in an extremely facile ring closure to acetylenylallene 116 and the isomeric 117. Subsequently, a solvent dependence study was conducted in which




of varying polarity. The results of this study are shown in Table VI.

TABLE VI
Solvent Effects Concerning the
Cyclization of 115

Solvent (5 ml/mole of <u>115</u>)	Acetylenyl Allene	
pyridine	1.5	1.5
acetylene chloride	1.5	1.5
chloroform	2.5	1.5
benzene	2.5	1.5
toluene	3.5	1.5

The great clearly indicates a marked preference for solvents of higher polarity, and, consequently, rules out a concerted C_4H_6 process. Special experiments of substituted cyclohexenes are known to be strongly solvent dependent and show the greatest increases in rate in those solvents which are better able to solvate the polycyclic intermediate.¹⁰ The data displayed in Table VI are consistent with this type of reaction.

could be obtained either by chromatography or by 100% air distillation. The results in some product decomposition and only slight purification. High resolution mass spectrometry of a distilled sample of 11 showed a parent ion at m/z 344 (M⁺) (calc for $C_{18}H_{20}N_2O_2$: 344.1480) and major peaks at m/z 334 and 332. A potential mechanistic pathway for ring




closure and fragmentation is illustrated in Scheme 12.

The stability 11 demonstrated in airway is useful as a facile dimer cyclization to give 12. This was taken out by the ability to survive vacuum distillation (by 85–100°C/0.75 mm Hg) without appreciable change.

Discussion

It was demonstrated in this chapter that the formation of quinoxalines and pyrazines from 2-aryl hydrazines under strong alkylation conditions was dependent upon the size of the substituents carried on the hetero nucleus of the hydrazine substrate. The systems with the greatest degree of bulk formed quinoxalines exclusively and exclusively while those with much smaller groups gave only pyrazines. In order to propose a meaningful mechanism which will effectively account for both products under the same conditions, one may consider further hypothetical factor. (Q) the preferential of diacylhydrazine-2-phenylhydrazide 10 was found to react rather with methyl dimethylformate or


SCHEME XX



loss of the proton analyzer results in the observed reaction 100. The proton source is methyl chloroacetate which arises from small amounts of hydrolysis which would be expected to produce methanol and chloroacetic acid. Bartley, Kuchinsk, and Reine²¹ have proposed a similar mechanism for the epimerization of dialkyl malonates to malonitriles (catalyzed by methylchloroacetic acid). The reluctance of ketonates 19 to undergo alkylation suggests a similar mechanism in the triaryl ketonate system. If the failure to observe alkylation is a function of electronics bound on nitrogen, and is probably so, then the triaryl system would, by virtue of the electronic withdrawing capacity of phenyl groups, tend to be even more reluctant. The proposed mechanism for the epimerization of *N*-acyl ketonates to malonitriles is illustrated in Scheme XII. The initial pathway in ketonates 100 could be an acid catalyzed process as shown. There is literature precedent for this type of behavior with diethyl carboallates which cyclize under acidic conditions to epimerizations.²¹ The nonspecificity of 100 relative to 81 would seem to be greatly enhanced by the resonance contribution of the nitrogen in the beginning. Alkylation at the *l*-position²² should take place easily to produce 82.

The comparative behavior of the *N*-acyl ketonate system in this work with those of other systems is interesting. The major product observed by Parker and Townsend¹⁰ from the thermolysis of dimethylacetone-*S*-phenylketone 20 was the epimerization 81. In addition, an uncharacterized isomer was identified by gas/ir as consisting approximately 10% of the product mixture. Subsequently, the identity of this isomer was not ascertained. The high percentage of dimer IPM 23, 80% in Parker's system is striking, since no trace of a substitution was detected when the

SCHEME 4304




with brominated **12** was treated at room temperature with methyl fluorosulfonate. Differences in experimental parameters such as solvent, base

acid concentration, solvent lot and temperature are expected to influence the course of reaction. In Barker's work, for example, the cyclization of benzimidazole were effected at 125° in a sealed tube without a solvent. Obviously, these differ considerably from the conditions described in this work. Such a dramatic increase in product simplification, though, may indicate a difference in activation energy for cyclization. If the energy of activation for cyclization is quite small as high with respect to chlorination, then temperature optimization would be expected to favor the latter as it did in this work. At higher temperatures, though, the two processes become competing and the thermodynamically more stable product (presumably the quinazoline 23) becomes dominant as it did in Barker's work. This hypothesis could be tested by simply conducting the alkylation of benzimidazole 21 with methyl chlorosulfonate at higher temperatures. The preponderance of a quinazolinone chlorosulfonate salt would confirm the suggestion.

In the case of triaryl benzimidazoles, the activation energy for chlorination may be low with respect to that for chlorosulfonation. This is conceivable if one compares the relative amounts of phenyl-phenyl populations existing in each species. Clearly, the quinazolinone suffers less from such intrusions than the corresponding triaryl. These arguments are summarized in Scheme XXXI.

The formation of quinazolinone from 2-alkyl benzimidazoles under alkylation conditions is more likely to be operating via direct methylation of the benzimidazole nitrogen. Support for this contention is found in the ease with which di-*n*-butylchloride-2-methylamine 22 (Scheme I) was alkylated with methyl chlorosulfonate. Also, a mechanism analogous to the one illustrated in Scheme XII has been proposed for

SCHEME 1811



SCHEME XXIV



The cyclization of 3-alkoxy-2-methyl-2-phenyl-1,3-dioxolanes to 3-alkoxy-2-methyl-2-phenyl-1,3-dioxanes (II) has also been observed by Harker and Ross⁶⁵ from the thermolysis of diphenylketene-3-methyl-2-phenyl-1,3-dioxolane. In a later paper they observed the reformation of 3-methyl-2-phenyl-1,3-dioxane from 3-methyl-2-phenyl-1,3-dioxolane under essentially the same thermolytic conditions. To this date, however, no attempts have been made concerning the mechanism of these thermolytic processes. For the most part, the product distribution of


via benzoylides, formed through thermal or photochemical cleavage of the benzoylides, which are known to be unstable under strong alkylating conditions. It is this author's belief that these acid anhydrides are responsible for all the heterocyclic diamine and diamine obtained under thermal conditions. The product could easily arise from a small amount of hydrogen chloride picked up by the bisoxazolone during synthesis, since all of the bisoxazolones used by Barker and Rowland¹⁰⁻¹² were prepared from the dehydrochlorination of imidized chloride precursors. Indeed, in two instances,^{10,11} respective diamine and/or triamine were isolated and characterized from bisoxazole distillates obtained during successive strip-up.

The epimerization of diisophtalimide-N-benzoylpyridine 22 to the corresponding isobutylene 23, could proceed by the mechanism already described in Scheme IIC. The failure to isolate or observe a triam from this system is probably indicative of rapid irreversible alkylation of the bisoxazole nitrogen. Once alkylation has taken place, ring closure to a triamine of the type described in this work is effectively inhibited.

ACKNOWLEDGMENTS


Introduction

Recent reports concerning the conversion of α -chlorosuccinimide to bisoxazolone isocyanatocarbonyl azides¹⁰⁻¹² has prompted this investigation into the synthesis of these systems in relatively hindered rings. Here, the potential C-halide and rather low (dimethylpyridine) It was hoped that deprotection could be effected by these bases, and provide a relatively straightforward route to bisoxazolone yields 22.




by trifluoride, however, took place much more slowly at room temperature, and even after 10 hours under refluxing toluene only ca. 5% conversion to **127** was obtained from our analysis of the crude product mixture.

The only other identifiable product was water **128** which crystallized with **127**. Further attempts at deprotection were made with the Eisteinmann salt **129**. Here, again only **127** and **128** were produced along with



expected small amounts of units III. The addition to III proceeded much more rapidly than with the *n*-chlorosuccinimide II to could be anticipated of a compound reacting on lactone carbon (11-12). It was hoped, however, that depolymerization would become a competing process. Unfortunately, there is no simple method for determining whether depolymerization is being achieved. Virtually all 1,3-dipolarophiles which could be used to interrupt the dipole would be unstable either to the strongly basic conditions needed for proper orientation¹² or to the electrophilicity of the bromination salt III.¹³⁻¹⁵ Qualitative experiments with N_2O would not serve well, since only a small steady state concentration of plus III is to be expected at any given time.¹⁶


Attention was focused on the synthesis of multifunctional units sterically protected at the oxygenation. Protection of this nature might discourage attack by base to form addition products similar to IX and XII, and perhaps direct the course of reaction towards depolymerization. Thus, the diethyl derivative XIII was prepared by vigorous treatment of units III with phosphorus pentachloride in refluxing benzene followed by dehydrochlorination with triethylamine. The *n*-chlorosuccinimide II



was obtained in 40% yield as a viscous yellow oil. This material proved to be very hygroscopic and again difficult to purify, however, preliminary data seemed to be the best technique for converting units IX to polymer. Treatment of *n*-chlorosuccinimide II with a benzene


slurry of sodium bis(trimethylsilyl)amide produced a deep reddish brown product mixture which showed no distinct infrared absorptions arising at 1028 cm^{-1} . This band is very characteristic of a heteroatom functionality and most likely arises from the desilylation of 111 to diglycidyl ether-*o*-methylated 10. Taylor and Smith¹⁸ have observed similar desilylations when treating various ketone bis(trimethylsilyl)amides with this same silylamide base.

Increasing the acidity of the protons carried by the aliphatic substituents was considered to be yet another alternative. The 3-benzyl-*o*-chloromandel 122 was prepared as a hygroscopic yellow liquid in 50% yield from its amide precursor 121. Because of inductive resonance effects of the phenyl group, the benzylic protons of 122 should be somewhat more acidic than those of the 3-methyl-. Consequently, α -



protons were abstracted under deprotonation conditions which could have been the direct result of initial proton abstraction. Indeed, only alcohols 124 and 125 along with smaller amounts of acids 126 were recovered from the product mixture. The allylamine product 123 was isolated in 50% yield while the vinylic derivative 127 was obtained as a mixture with acids 128 in the approximate molar ratio of 2:2:1 respectively.


In a final effort to convert an *o*-chloromandel with appropriately acidic protons adjacent to nitrogen, the 3-(2-chloroethyl) amide 129



was synthesized via the condensation of ethyl 9-oxoethylacetate 127 with isobutyryl chloride 128. The conversion of 127 to the α -chloro-acetate 129 was effected by the usual methods, but only in ca. 10% yield.




This hygroscopic colorless liquid exhibited infrared bands at 1755 and 1660 cm^{-1} corresponding to stretching modes of the ester carbonyl and ketone functionalities respectively. The C-methyl maximum of the acetate appeared as sharp singlets at δ 1.80 and δ 1.85 in carbon tetrachloride, but broadened seriously when dissolved in chloroform. In addition to the expected singlet pattern at δ 3.23 (singlet) and δ 4.30 (quartet), 9-methyl and 9-oxobutano singlets were



observed at δ 3.47 and δ 3.17 respectively. When the *n*-chlorosuccinimide **127** was treated with potassium *n*-butyrate in acetonitrile ether for 12 hours under a nitrogen atmosphere, a mixture was obtained whose gross analytical properties were essentially identical to those of acid **128**. Other resonances appeared at δ 3.63 (doublet, CH_2 -OH), δ 4.51 (singlet, CH_2 -OH) and 5.62 (singlet, CH_2 -OH), but represented only a small amount (5%) of the total product mixture. These signals could correspond to the




isopropylsuccinimide **129**. Due to the large amount of contamination, and the small amount of material at hand, attempts to isolate this compound were not undertaken. The *n*-chlorosuccinimide **127** exhibited an interesting property when left in solution with methylene chloride or chloroform. After extended periods of time in these solvents, cyclization to the succinimide salt **131** occurred almost quantitatively. When isolated **131** was spectroscopically, the signals associated with **127** gradually disappeared, and, finally, were completely replaced by resonances



at δ 1.40 (heptet, $J=7$ Hz), 1.22 (triplet, $J=7$ Hz), 3.88 (multiplet, $J=7$ Hz), 4.17 (septet), 5.44 (quartet), and 8.25 (singlet). Accurate integration of these signals was not possible due to the overlapping of residual acids 131 resonances present as a result of hydrolysis of 130. In order to unequivocally establish the structure of 131, the 3-isobutyranal 132 was prepared by converting the acid 131 to its benzoyl anhydride and subsequent dehydrochlorination with triethylamine. This

SCHEME XXV




compound was identical in every respect to that reported in the literature by another author.⁷⁶ Isomerization of 115 to isopropenyl vinyl ether with methyl fluorosulfonate was efficient, smooth and in high yield. This isopropenyl substituted an infrared absorption band at 1655 cm^{-1} and one signal at $\delta\ 3.44$ (doublet, $J=2\text{ Hz}$, $2H$), 3.47 (doublet, $J=2\text{ Hz}$, $2H$), 3.12 (quartet, $J=7\text{ Hz}$, $2H$), 3.88 (singlet, $2H$), 4.38 (quartet, $J=7\text{ Hz}$, $2H$), and 5.13 (singlet, $2H$). The positions of these signals are quite similar to those of 111 with the exception of the vinyl protons which appear at the very low value of $\delta\ 3.12$. No rational explanation can be offered for this difference in the chemical shifts of the vinyl protons. It is difficult to ascribe the magnitude of this difference merely to a difference in conformation, but when this idea too appears to be identical. Furthermore, sodium borohydride reduction of 111 and 115 give the same 3-methyl-2-butethylalcohols 153.

The bicyclic 116 is of interest, too. Simple decomposition of this molecule will lead to a product discussed earlier and thought to be 112 (vide supra). The characteristic behavior of 116 has not been investigated. However, it appears to have the potential of reacting itself as a Cornforth type⁷⁷ ring opening in the ketone-alcohol yield, and, possibly, closer to the isopropylidene solution 114.

Discussion

The ability of *n*-chlorosuccinates as promoters in ketonolysis yields 52 is linked to several reactions: (i) the ease with which nucleophilic displacement occurs at the vinyl carbon in these esters under decomposition conditions, (ii) the difficulties encountered in synthesizing variously protected derivatives, and (iii) the apparent low ability of the proposed adjacent to arrange. The facile displacement

SCHEME XIV




of chloride at the vinyl carbon by a sterically hindered nucleophilic base was surprising in view of the fact that vinyl substrates are normally very resistant toward substitution.²⁸ This unusual observation is explainable in terms of a pre-ionization in the bromination with 12 as illustrated in Scheme XVII. This pre-ionization mechanism could also explain the cyclization of 12 to its monocyclic chloride salt 15. Interestingly, this cyclization proceeds in acrylonitrile or chloroform, but not in other polar solvents such as acetonitrile or dimethylformamide.

SCHEME XXVII




from the addition of hydrogen chloride which is known to be present in chloroformal solvents failed to induce cyclization. Apparently, other differences in these solvents are responsible for this rather novel transformation. In any case, more work is required to determine the subtle effects of solvent media upon cyclization.

SCHEME XXVIII



The large amount of water 12 detected in the product films, resulting from the accepted decomposition of 12 with protic or α -hydroxide was precluding besides the use of anhydrous solvents and efforts to maintain a moisture free environment. It is true that atmospheric moisture could have been responsible for its formation, however, one cannot overlook the possibility of a decomposition mechanism illustrated in Scheme XXIX.⁸

SCHEME XXIX




CHAPTER 102


REACTIONS OF MONOTERPENOIDS IN ALIPHATICS

Cyclizations involving six-membered transition states are favored by the formation of closure of two sigma bonds involving six π single bonds (Scheme XXX).⁷³ Further examples include small molecule

SCHEME XXX



reactions and various as well as other studies to clarify. Electrophilic cyclizations of a 34 type are extremely rare.⁷⁷ The most recent report of a 34 cyclization involved the addition of a 4-hydroxy-1,5-diene to the bicyclic compound 11 with formation of the bicyclic diol 12.⁷⁸ Although the reaction almost certainly proceeded 12 via the 1,5-dipolar intermediate bicyclic intermediate, no conclusions could be drawn concerning the mechanism of the addition of the diene to the alkyne. Since the properties of 11 and 12 did not lead themselves to a detailed mechanistic study, systems were sought after which were likely to undergo such cyclizations and which might be mechanistically informative.



Functionally substituted naphthalenes have often been used to determine relative preferences with respect to their ability to undergo thermal or photochemical ring opening to their 1,3-dicarbonyl isomers. Also, cycloadditions of naphthalenes with various 1,3-dicarbonyls are known to take place in a concerted fashion,⁴¹⁻⁴³ and have been shown to conform with orbital symmetry requirements advanced by Woodward and Hoffmann.⁴⁴

Assuming a Diels adduct is formed between a 1,1,2-trisubstituted naphthalene and *n*-butylmaleimide 1, one could expect to find two possible adducts (189 and/or 190). The mechanistic pathway, of course, will dictate the product distribution. This will be dependent upon several



189



190

known: (1) a concerted sig, 6 stepwise addition, (2) the configuration of the sig, trans of the addition,^{11,12} (3) the rate of epoxidation relative to that of addition decomposition,^{13,14} and (4) the experimental conditions.

The functionalized alkydine system was chosen because ring opening to the addition yield has been demonstrated to be clean, reversible, and not complicated by troublesome side reactions. Furthermore, the small number of protons on the alkydine ring (and eventually on 11f and 12f) would allow easy access to mechanistic studies sig and spectroscopy.

Results

The alkydine chosen in this work for the purpose of generating a 1,3-dipolar intermediate was ethyl trans-4,5-dihydro-3-methylene-oxaphosphate 11f. Its synthesis was accomplished according to equation (VII) by an already published procedure.¹⁵ Alkydine 11f was shown to



equilibrate easily to rotating carbon tetrachloride to an approximate 65:35 sig trans isomer distribution. Treatment of alkydine 11f with *o*-hydroquinoline 1 in a sealed tube for 24 hours at 80° dispropagately produced an adduct with the formula 12f. However, spectroscopic investigation of the reaction mixture showed that the alkydine had indeed isomerized to a sig trans mixture. In view of this, we concluded that the unreacted 11f had been formed and failed to react.



Unexpectedly, chromatography of 122 on H^+ in a carbon tetrachloride solution containing a twofold excess of *n*-butylmagnesium bromide furnished 123 as a white oil in 70% yield.⁽⁴⁷⁾ Repeated attempts to



crystallize 123 resulted in failure. Purification by column chromatography afforded a sample of near analytical quality which was subsequently used for characterization and structure elucidation. Evidence for the assigned structure 123 came from the mass spectrum which shows a parent ion at m/e 180. The infrared spectrum displayed absorption bands at 3000 and 1740 cm^{-1} corresponding to the ketone and ester functionalities respectively. The nmr spectrum included a set of two singlets at δ 4.16 superimposed over a few peaks quartet centered at δ 4.15. These data, however, do not unequivocally rule out the possibility of the isomeric ketone 124.

Spectral properties of acyclic acid compounds in the area of chemical shift data of the methylene protons of 121 and 122 suggested

**150****151**

Leaves **151** as the more plausible structure. It might be expected, however, that the inductive effect of the hydroxyl group would be in competition with the steric lone pair in both structures **150**.

**152****153**

$$\text{H}_\text{a} = \delta \quad 3.7\delta$$

$$\text{H}_\text{b} = \quad 4.1\delta$$

$$\text{D}_\text{H} \quad \text{C}=\text{O} = 1730 \text{ cm}^{-1}$$

$$\text{H}_\text{a} = \delta \quad 4.1\delta$$

and **152** should have a slight deshielding effect on the methylene protons in either case. If this assumption is correct, model compound **151** would be more compatible with the experimental data, and thereby be in agreement with the assigned structure **151**.


Further and multiproton structural confirmation was obtained from the acid-catalyzed hydrolysis of lactonide **155** to yield **154**, and the subsequent synthesis of **154** from 2-*trans*-4-phenoxy-5-*γ*-butyrolactonide **156** and ethyl 4-phenoxybutyrate **157** as outlined in Scheme 100. The

Discussion

The origin of intermediate 120 can be rationalized on the basis of two mechanistic proposals. The first, illustrated in Scheme VIII, as mechanism 1, involves the reversible ring opening of anhydride 123 to 124. Attack by vinylbenziodole in a concerted addition would most likely occur at the apparently more electrophilic benzoyl carbonyl of the anhydride plus 124 and lead to the polymerically inactive adduct 125. Typically substituted anhydrides have been noted to show analogous

SCHEME VIII


Mechanism 1.



isolated when thermolyzed in the presence of nucleophilic reagents.^{11,12} Intramolecular proton transfer to the ortho-chloromethyl solvent moieties¹³ results in the direct formation of product 115. Mechanism A, electrophilic attack, suffers from the observation that initial attempts to promote [W] addition by the thermolysis of aziridine 111 with excess isocyanide in the absence of a solvent resulted only in the isomerization of the aziridine. If the aziridine dipole was the sole species for adduct formation, mechanism A would be in agreement with experimental facts. Apparently, this is not the case. The properties of the solvent (carbon tetrachloride) must in some fashion be related to the overall transformation. Mechanism B accounts nicely for the catalytic activity afforded by the solvent. Protonic coupling of the initially formed


SCHEME XXXIII

Mechanism B



yield 18, produces the addition only 18, which could be susceptible toward nucleophilic attack by the isocyanide to give 19. Loss of a proton results in isocyanide 21 and regeneration of the acid catalyst.

The effects of trace amounts of gold in chlorinated solvents such as carbon tetrachloride and chloroform have been demonstrated by Hill,³⁴ Yee,³⁵ and others³⁶ to be responsible for isocyanides obtained from the treatment of selected ketone isomers with *n*-butylisocyanide 2 in these solvents. The capture of protons by isocyanides was shown by Harris³⁷ to occur readily with functionalized isocyanides of the type used in this work. And, when carried out in the presence of a variety of nucleophiles, addition products 22 were isolated in yields ranging from 20-90%.



The question of why *tert*-butylisocyanide produces between *n*-butylisocyanide 2 and isocyanide 21 more (or decreases still remains pending. The lack of correlation between the isocyanide label and isocyanide yield systems in this respect is not immediately obvious. It is apparent that more nucleophile may need to be used in this case.

CHAPTER IV

EXPERIMENTAL

Boiling points and melting points are recorded to degrees Celsius and are uncorrected. Boiling points were determined with a Thomas-Hoover Schmidt auxiliary swirling joint apparatus using 1-4-1.6 x 70 mm glass capillary tubes. Boiling points were determined by conventional distillation techniques or by micro capillary methods. Infrared spectra were recorded on a Perkin-Elmer Model 577 Infracolor Prism spectrometer, and collected at 3000 cm⁻¹ with a polyethylene film. Raman spectra were recorded on a Perkin-Elmer Model 560-R analytical spectrometer for 4000 cm⁻¹ region spectra, and a Perkin-Elmer Model 521-R spectrometer for 1600 cm⁻¹ region and proton decoupled spectra. Carbon-13 nuclear magnetic resonance spectra were obtained from a Varian Model XL-100 spectrometer operating at a probe frequency of 125.6 MHz. All chemical shifts (δ) in designated solvents are reported in parts per million (ppm) downfield from tetramethylsilane as an internal standard. Besides low resolution mass spectra, exact mass and molecular weight data were measured on an AEI MS-90 double beam spectrometer at an ionizing potential of 10 eV. Microanalyses were carried out by Arminio Marchini, Inc., Atlanta, Georgia, and in all cases are in agreement with assigned structures. NMR spectra, reactions and manipulations of sensitive materials were carried out in a laboratory fume hood with a continuous stream of nitrogen. Solvent evaporation was performed at reduced pressure on a Buchi Rotavapor-R rotary evaporator equipped with a water aspirator.

1-Benzyloxycarbonyl Chloride (28)

1-Benzyloxycarbonyl chloride was prepared from 1-butyl chloride and tripropylamine according to the procedure described by Milne and Koberstein,¹⁷ and standardized according to the method of Gilman, *op. cit.*¹⁸

2,2,3,3-Tetrachloro-1,3-Propanediol (29)

2,2,3,3-Tetrachloro-1,3-propanediol was prepared from tripropylamine chloride and phosgene chloride with cuprous chloride as a catalyst according to the procedure described by East and Fawcett¹⁴ by 152° in liquid phase (445 cm^{-1} C=O, nm CHCl_3 δ 1.13 (singlet, 3H).

1,2-Bis-(4-benzyloxy) Chloride (30)

The synthesis of 1,2-bis-(4-benzyloxy) through 1,2-di-(4-benzyloxy) chloride were carried out according to the methods described by Brown, Whittell and Roberts.¹⁵ The product was obtained as a colorless liquid bp $10-10^\circ/12$ mm Hg in liquid phase 1700 cm^{-1} C=O; nm CHCl_3 δ 1.12 (singlet, 3H), 1.48 (singlet, 3H).

2,2-Diphenyl-3,3-Dichloropropanediol (31)

Acetylene ethylenes (25.0 g, 0.33 mole) was placed in a 2 x 30 cm Barker and Porter flask (also previously cooled to 0°) with an ice water bath. 1,2-Di-(4-benzyloxy) chloride (2.31 g, 44.8 mmole) was added directly to the chilled ethylenes without stirring. The flask was sealed, removed from the ice water bath, shaken to effect mixing of reagents, then positioned over a magnetic stirrer where the mixture was stirred for 16 hours. The gummy product mixture was

current ethylenes. The residue was again taken up in chloroform, washed with dilute hydrochloric acid, then dried over anhydrous magnesium sulfate. Evaporation of the solvent left a pale yellow solid. Recrystallization from hexane afforded 2-methyl-2,3-dichlorobutanoic acid (1.7 g, 30%) as colorless prisms mp 47-48°C, lit (14c) 48-50 (decolor), strong IR (KBr, 1740 cm^{-1} (C=O, weak), and (CHCl₃) δ 1.20 (singlet, 3H), 1.12 (triplet, 3H), 3.0, 3.0, 3.45 (quartet, 2H), 3.12 (quartet, 3H), 3.25 (quintet, 1H).

ANAL. Calcd for C₄H₇Cl₂O: C, 42.34; H, 33.56; O, 24.10.

Found: C, 42.34; H, 33.43; O, 24.0.

2,3-Dichlorobutanoic acid (15)

A solution of 2-methyl-2,3-dichlorobutanoic acid (1.35 g, 5.4 mmol) in 15 ml of benzene was prepared in a 100 ml round-bottomed flask equipped with a reflux condenser and a magnetic stirring bar. Triphenyl phosphorite (2.29 g, 5.4 mmol) was added and the resulting suspension refluxed for approximately 30 minutes. Benzene and phosphoryl chloride were removed by evaporation at reduced pressure, and the resulting residue of phosphoryl chloride added with 10 ml of benzene. The crude 2-methyl chloride was treated directly with triethylamine (3.32 g, 2.74 ml, 21.8 mmol) in 30 ml of benzene, and the resulting mixture refluxed for 1 hour. The resulting triethylamine hydrochloride was removed by vacuum filtration and the filtrate concentrated at reduced pressure to a brown liquid. Vacuum distillation (Baird-Porter, 45-50°/0.25 mm Hg) gave 2,3-dichlorobutanoic acid (1.54 g) concentrated to the degree of approximately 100 by which appears to be 2,3-dichlorobutanoic acid¹⁰ IR (liquid film) 1740 cm^{-1} (C=O); and (CHCl₃) δ 1.01 (singlet, 3H), 1.12 (triplet, 3H), 3.0 (quartet, 3H), 3.12 (quintet, 3H); n_D²⁰ 1.444.

for $C_{11}H_{11}NO$ 181.1458, (found) 181.1402.

Diethylhexamethylenediphenylamine Fluorosulfonate (20)

Freshly prepared diethylhexamethylenediphenylamine (1) 40 g was placed into the septinuder bulb into a 100 ml round-bottomed flask with 50 ml of anhydrous ether. The flask was then set up for magnetic stirring, and protected from atmospheric moisture with a calcium sulfate drying tube. Diethyl fluorosulfonate (1) 31 g, 1.00 ml, 13.1 mmol was syringed into the reaction mixture, and the resulting mixture allowed at room temperature for approximately 30 minutes. Precipitation of the hexamethylenediphenylamine salts as a white fluorosulfonate suspension took place within moments after the introduction of the diethyl fluorosulfonate. The colorless solid was collected on a small Hirsch funnel by vacuum filtration and washed several times with anhydrous ether. Recrystallized from anhydrous diethyl ether afforded di-*n*-butylhexamethylenediphenyl-*n*-butyl-*n*-ethylamine fluorosulfonate (2) 30 g as a white powder mp 154-155° (decaying), d_4^{20} 0.91 (20°) 0.909 cm^{-1} (20°), n_D^{20} 1.411 (20°) (singlet, 3H), 1.48 (triplet, 3x3 H), 3.78 (singlet, 3H), 4.11 (quartet, 3x2 H), 5.9.

n-Butyl-*n*-Ethyl-1,1,2,2-tetrafluoroethane (21)

Di-*n*-butylhexamethylenediphenyl-*n*-butyl-*n*-ethylamine fluorosulfonate (2) 30 g, 0.14 mmol and 15 ml of distilled water were placed in a 10 ml triaxial flask. The solution was made basic by adding 5 ml of 50% aqueous sodium hydroxide, and the resulting mixture allowed at room temperature for approximately 18 hours. Extraction into three 15 ml portions of diethyl ether followed by drying over anhydrous magnesium sulfate and evaporation at reduced pressure produced *n*-butyl-*n*-ethyl-

Treatment of 2,4,6-Trichloro-3-(2-methyl-5-oxopentylidene)-1,3,5-triazine-1,3,5-trioxide with Diethylphosphorylbenzide to Form Diethylphosphorylbenzide (1975)

A solution of sodium bis(trimethylsilyl)benzide (6.495 g, 3.44 mmol) in 50 ml of dry hexamethylphosphoramide was prepared in a 50 ml round-bottomed flask equipped with a magnetic stirring bar and a rubber septum. A solution of diethylphosphoro-3-methyl-5-oxopentylidene diacetate (5.445 g, 1.41 mmol) in 1 ml of dry hexamethylphosphoramide was syringed into the stirred benzene solution. The resulting homogeneous mixture was stirred at room temperature for 24 hours. The mixture was diluted with 50 ml of water, then extracted four times with 50 ml portions of diethyl ether. The ether solutions were combined, concentrated at reduced pressure, and vacuum distilled (Bp: 100°C) to remove the remaining hexamethylphosphoramide. The residue was taken up in ethylacetone, which, after several hours of refrigeration, produced glyoxal (1975) (5.445 g) in 21.1% yield.

The distillate was taken up in 50 ml of diethyl ether, washed three times with 15 ml portions of water, dried over anhydrous magnesium sulfate, and evaporated at reduced pressure to a pale yellow liquid (5.445 g). This material showed major mass signals at m/z 100 (base), 140, 1.36 (mole), 1.67 (m, 10), 1.55 (mole), 5.740, 3.8 (mole), 1.250, and 1.34 (mole), 1.67 (m, 10) and was shown by comparison with an authentic sample to be 3-methyl-5-oxopentylidene-2,4,6-trichloro-1,3,5-triazine (1975).

Run 8-ethyl-1-(1,2,4,4-tetramethyl-3-oxocyclopentylidene)pyrrolidine, 7.
Reagents: 1-(1,2,4,4-tetramethyl-3-oxocyclopentylidene)-2-pyrrolidinyl
1,2,4,4-tetramethyl-3-oxocyclopentane, 10.0 g, 0.040 mole; 2-pyrrolidinyl
1,2,4,4-tetramethyl-3-oxocyclopentane, 10.0 g, 0.040 mole; 2-pyrrolidinyl
1,2,4,4-tetramethyl-3-oxocyclopentane, 10.0 g, 0.040 mole.

2,2-bis(4-ethyl-1-(1,2,4,4-tetramethyl-3-oxocyclopentylidene)pyrrolidinyl)propane

26.600 g, 0.080 mol), methanol (2.10 g, 12.8 mol), and 10 ml of dry toluene were combined in a 30 ml round-bottom flask equipped with a magnetic stirring bar and a condenser cooled with dry ice. The flask was transferred to a drybox and positioned over a magnetic stirrer where sodium bis(trimethylsilyl)acetate (6.75 g, 0.080 mol) was added. The resulting suspension was then cooled to room temperature under nitrogen atmosphere for 12 hours. The golden granular mixture was poured through a dilute funnel containing a bed of Dillite, and the pale yellow fragments at reduced pressure to remove solvent and excess methanol. After removing the residual traces of solvent by azeotropic distillation at high vacuum, 2,2-bis(4-ethyl-1-(1,2,4,4-tetramethyl-3-oxocyclopentylidene)pyrrolidinyl)propane 10.344 g, 44.50 mmol was obtained as an solid semicrystalline, pale yellow with $n_D^{20} 1.4803_{10} (0.1) \text{ cm}^{-1}$ ($d_4^{20} 0.970$), $n_D^{25} 1.4783_{10}$ (triplet, 1-2 Hz, multiplet), 1.20 (singlet, 1-buty), 1.30 (singlet, 1-buty), 1.2-1.8 (multiplet, methylene and methine), ^{13}C see $^{13}\text{C}(\text{CDCl}_3)$ 15.8 (quartet, $-\text{CH}_2-\text{CH}_2-$), 26.2 (triplet, C_β), 31.1 (quartet, 1-buty), 31.1 (triplet, C_β), 31.5 (quartet, 1-buty), 32.8 (triplet, 1-buty), 36.8 (triplet, 1-buty), 36.7 (doublet, C_α), 41.2 (quartet, C_α), 41.5 (triplet, C_β), 46.2 (triplet, $-\text{CH}_2-\text{CH}_2-$), 51.1 (doublet, C_{CH_3}), 54.4 (doublet, C_{CH_3}), 57.4 (quartet, C_α), 174.8 (triplet, carbonyl C=O), 181.4 (triplet, carbonyl C=O), n/a (quadrupole $\text{C}_{11}\text{H}_{20}\text{N}_2\text{O}$ DBP, 1740, 1740, 1740, 1740).

The crude lactone, was diluted with approximately 5 ml of benzene and applied to a neutral alumina column (1.5 x 30 cm) packed in pyridine vapor (150°C). The 40 ml fractions were collected with

Dimer, $C_{12}H_{16}$, 136.8 (singlet, C_2), 131.2 (singlet, C_7H), 142.5 (singlet, $C_{10}H$), 145.8 (singlet, $C_{12}H$), 145.8 (singlet, $C_{12}H$), 148.2 (singlet, $C_{10}H$), 155.7 (singlet, C_2); *Qucid* for $C_{12}H_{16}O$ 155.2145, (Found) 155.1801.

1,1-Dichloroacetyl Chloride

1,1-Dichloroacetyl chloride (178.8 g, 6.125 mole) and thionyl chloride (147.8 g, 67.5 ml, 4.26 mole) were combined in a 100 ml round-bottomed flask equipped with a magnetic stirring bar and a reflux condenser. The slurry was refluxed for 4-5 hours, during which time the mixture became homogeneous and developed a yellow appearance. The excess thionyl chloride was removed by evaporation at reduced pressure and the residue vacuum distilled at 123°/5.5 mm Hg (lit.¹⁰ bp 170-171°/16 mm Hg). 1,1-Dichloroacetyl chloride (178.2 g, 67.125) was obtained as a yellow-orange liquid which solidified on standing. ^1H NMR ($CDCl_3$) 1.75 (m¹¹ 4H-4H) and 5.23 (singlet, 2H), 7.18 (singlet, 2H).

N-Phenyl-2,1-bisphosphorylamide (II)

A solution consisting of sodium (14.8 g, 0.625 mole), triethylamine (114.8 g, 104. ml, 0.625 mole) and 200 ml of benzene was prepared in a 1000 ml three-necked round-bottomed flask equipped with an addition funnel, mechanical stirrer, and an ice water cooling bath. 1,1-Dichloroacetyl chloride (178.2 g, 6.125 mole) in 100 ml of benzene was added dropwise to the above sodium solution and the mixture stirred for approximately 18 hours. The triethylamine hydrochloride was collected by sodium filtration and washed several times with 100

¹¹Monomer signals thought to be due to minor $(E)-CH_2$ isomer.

aniline hydrochloride. The colorless solid which remained was collected and dried. The benzene filtrate was washed with dilute hydrochloric acid, dried over anhydrous magnesium sulfate and evaporated to give a second crop of colorless solid. The two crops were combined and recrystallized from benzene/petroleum ether (60-120°). *N*-Phenyl-2,2'-diphenylacetamide 126.8 g, 58% was isolated as a fluffy colorless solid mp 176.5-180.2° (lit.²⁰ mp 180°); λ_{max} 283m μ 2445 cm⁻¹ (C=O), ν_{max} 3361 ν 4 3-45 (strong), 30-, 1,10-3-55 (weak), 158 ν .

Diethylketone-*N*-Phenylamine (17)

Diethylketone-*N*-phenylamine was prepared by aniline iodide dehydrohalide of the *N*-dimethylbenzyl chloride procedure according to the procedure described by Norrish and French²¹ mp 52-53° (lit.²¹ mp 55-56°), λ_{max} 283m μ 1798 cm⁻¹ 3000-3100, ν_{max} 3361 ν 4 3-45 (weak), 30-, 1,10-3-55 (weak), 158 ν .

Synthesis of Diphenylacetone-*N*-Phenylamine (18)

A solution of diphenylacetone-*N*-phenylamine (1.50 g, 4.36 mmol), and 25 ml of anhydrous ether was prepared in a 50 ml round-bottomed flask equipped with a magnetic stirring bar and a calcium chloride drying tube. This solution was treated with methyl fluorosulfonate (6.50 ml, 4.76 g, 4.36 mmol) and stirred at room temperature for a period of 48 hours. The clear colored quinazolinone salt (1.30 g, 84%) was isolated in a small Erlenmeyer flask and washed several times with ether to remove excess methyl fluorosulfonate and excess pyridine. Purification was achieved by dissolving the crude fluorosulfonate salt in a minimum amount of hot dichloroform and diluting with approximately

260 ml of hexane. The resulting solution produced brownish granules of 1-methyl-2-methylcyclo-3-phenyl-4-benzoylpyridine-3,4-dihydropyrimidin-2-one dihydrochloride after several hours of refrigeration; mp 145-154° (decorp.); n_D (c=0.5mm, CHCl₃) = 1.58 (singlet, 3H), 5.1 (singlet, 1H), 6.4-6.7 (multiplet, 2H).

1-Methyl-2-phenyl-3-phenyl-4-benzoylpyridine-3,4-dihydropyrimidin-2-one 180

1-Methyl-2-benzoylpyrid-3-phenyl-4-benzoylpyridine-3,4-dihydropyrimidin-2-one dihydrochloride (1.88 g, 4.15 mmol), 25 ml of 1,4-dioxane, and 50 ml of 2N sodium acetate were combined in a 250 ml three-neck flask and stirred magnetically until a clear yellow solution resulted (20, 30 minutes). The reaction mixture was placed on a rotary evaporator where most of the water and dioxane were removed at reduced pressure. The sticky yellow residue which remained was taken up in chloroform, dried over anhydrous magnesium sulfate, then evaporated to dryness. Recrystallization from petroleum ether (45-140°) produced 1-methyl-2-phenyl-3-phenyl-4-benzoylpyridine-3,4-dihydropyrimidin-2-one (1.5 g, 64%) as a transparent yellow solid, mp 200-204.5°; n_D (CHCl₃) = 1.58 (singlet, 3H), 6.4-6.7 (multiplet, 2H), n_D 1.5885 (for C₂₁H₁₄N₂O) 231.2364, $\lambda_{\text{max}}^{\text{CHCl}_3}$ 242.2181.

1-Methyl-2-benzoylpyrid-3-phenyl-4-benzoylpyridine-3,4-dihydropyrimidin-2-one 192

1-Methyl-2-benzoylpyrid-3-phenyl-4-benzoylpyridine-3,4-dihydropyrimidin-2-one dihydrochloride (1.88 g, 4.15 mmol) and 25 ml of ethanol were combined in a 250 ml three-neck flask equipped with a magnetic stirring bar. Sodium hydride (8.25 g, 3.74 mmol) was added in several small increments to the stirred solid solution. After the

Crystallization of the triphenyl ether gave red ECD obtained
 product 1,1'-diphenyl-2-(4-*B*-methoxyphenyl-2-diphenylacetylphenyl)
 ethylene (2.114) g, 1400 in thin brown yellow needles mp 162.3-164°
 in CHCl_3 1300 (s-ol), 1405 cm^{-1} C=O, ν_{max} 3300 (s) 4.28 (doublet,
 J=8 Hz, 2H, collapse to a singlet with $\delta_{\text{H}}^{\text{DMS}}$ 1-4) (singlet, 2H,
 6.45 (doublet, $J_{\text{H}}^{\text{DMS}}$ 8.8 Hz, 2H, $\delta_{\text{H}}^{\text{DMS}}$ 6.23 (singlet, 1H), 4.4-7.4
 (multiplet, 2H), 7.17 (doublet, $J_{\text{H}}^{\text{DMS}}$ 8.8 Hz, 2H, $\delta_{\text{H}}^{\text{DMS}}$ 8.38 (broad, 2H,
 water out with $\delta_{\text{H}}^{\text{DMS}}$, ν_{max} 3300 for $\text{C}_{11}\text{H}_{13}\text{NO}$ 178-180, 1190-1210.

The filtrate from the above crystallization was concentrated at
 reduced pressure and applied to a silica gel column packed in carbon
 tetrachloride. The final fraction eluted with 40 ml of carbon
 tetrachloride contained an unidentified material not now discussed
 a second fraction was collected with 80 ml of methylene chloride
 carbon tetrachloride (9:1), and concentrated at reduced pressure to a
 yellow oil (2.170 g). Spectroscopic (see) analysis of this compound
 indicated the presence of 8-methyl-2-*B*-diphenylacetylphenyl (2.115) g,
 210. An analytical sample was prepared by vacuum distillation
 (b.p. 178-179°/0.2 mm Hg) in CHCl_3 1300 (s-ol), 1410 cm^{-1}
 C=O, ν_{max} 3300 (s) 4.28 (doublet, J=8 Hz, 2H, collapse to a singlet with
 $\delta_{\text{H}}^{\text{DMS}}$ 1-4) (singlet, 2H, 6.23-6.45 (multiplet, 2H, $\delta_{\text{H}}^{\text{DMS}}$ 6.23,
 6.45 (doublet, 1H), 7.18 (doublet, $J_{\text{H}}^{\text{DMS}}$ 8.8 Hz, $J_{\text{H}}^{\text{DMS}}$ 1.5 Hz, 2H, $\delta_{\text{H}}^{\text{DMS}}$
 8.45 (broad, 2H, water out with $\delta_{\text{H}}^{\text{DMS}}$, ν_{max} 3300 for $\text{C}_{11}\text{H}_{13}\text{NO}$
 181-184, 1190-1210.

Identification was based on spectroscopic comparison with an
 authentic sample.

N-(p-Tolyl)-2,3-Diisopropylsuccinimide (31)

A solution consisting of pyridine 68.4 g, 0.415 mole), triethylamine (25.4 g, 0.415 mole), and 500 ml of benzene was prepared in a 1000 ml three-necked round-bottomed flask equipped with an addition funnel, mechanical stirrer, and an air-water cooling bath. 1,2-Diisopropylsuccinyl chloride (26.4 g, 0.115 mole) in 100 ml of benzene was added dropwise to the above color solution and allowed for 3 hours. The triethylamine hydrochloride was collected by aqueous filtration and washed several times with 10% aqueous sodium hydroxide. The colorless solid which remained was collected and dried. The benzene filtrate was washed with dilute hydrochloric acid, dried over anhydrous magnesium sulfate then concentrated in a second crop of product. The two crops were combined and recrystallized from benzene/petroleum ether. N-(p-Tolyl)-2,3-Diisopropylsuccinimide (26.4 g, 33%) was isolated as a colorless solid mp 171-176.5° C(lit.⁷⁰ mp 175-177°), IR (KBr)₂ 1675 cm⁻¹ (C=O), n_D²⁰ (CHCl₃) 1.42 (singlet, 6H, 3-CH₃ doublet, 4H, 4.00-4.25 (multiplet, 16H)

Diisopropylsuccinimide-N-(p-Tolyl)Imide (32)

Diisopropylsuccinimide-N-(p-tolyl)Imide was prepared by the direct dehydration of the corresponding acid according to the method described by Korman and Knight¹⁴ mp 41.5-45.5° C(lit.¹⁴ mp 45-46°); IR (KBr) 1745 cm⁻¹ (C=O); n_D²⁰ (CHCl₃) 1.42 (doublet, 6H, 4.50-4.67 (multiplet, 16H).

Acetylide of Diisopropylsuccinimide-N-(p-Tolyl)Imide (33)

A solution of diisopropylsuccinimide-N-(p-tolyl)Imide (8.00 g, 16.8 mmole) and 25 ml of anhydrous ether was prepared in a 100 ml round-bottomed

flask equipped with a magnetic stirring bar and a reflux condenser drying tube. This solution was treated with methyl fluorosulfonate (5.00 ml, 5.00 g, 22.2 mmol) and stirred at room temperature for 48 hours. Precipitation of the crude polysulfonate salt occurred suddenly after the first 14 hours of stirring. In some instances, cooling or scratching with a glass rod was necessary to induce crystallization. The free sulfonate 1,4-bis(2-hydroxy-2-propyl-5-benzoylphenyl)-2-(p-tolyl-5-benzoylphenylidene)-3,4-dihydroquinazolinium fluorosulfonate (4.40 g, 840) was collected in a small filter funnel and washed several times with anhydrous ether to remove excess methyl fluorosulfonate and other byproducts. Purification ggs crystallization was achieved by the addition of 110-120 ml of hot benzene to a solution of the crude polysulfonate salt in 20-25 ml of dichloromethane. Ten platelets of 1,4-bis(2-hydroxy-2-propyl-5-benzoylphenyl)-2-(p-tolyl-5-benzoylphenylidene)-3,4-dihydroquinazolinium fluorosulfonate appeared on cooling up 340-348° (decap-3) and (trifluoroacetic acid) 5 2.15 (singlet, 3H), 4.33 (singlet, 3H), 5.34 (singlet, 3H), 6.3 (singlet, 3H), 6.38-7.14 (multiplet, 17H).

1,4-Bis(2-hydroxy-2-propyl-5-benzoylphenyl)-2-(p-tolyl-5-benzoylphenylidene)-3,4-dihydroquinazolinium
(mg, 100)

1,4-bis(2-hydroxy-2-propyl-5-benzoylphenyl)-2-(p-tolyl-5-benzoylphenylidene)-3,4-dihydroquinazolinium fluorosulfonate (5.00 g, 2.22 mmol), 25 ml of 1,4-dioxane, and 20 ml of 32% sodium hydride were combined in a 100 ml nitrogen-flask and heated in a steam bath for 30 minutes with intermittent stirring. The reaction mixture was placed on a rotary evaporator where most of the water and dioxane were removed at reduced pressure. The gummy yellow residue which remained was taken up in dichloromethane, dried over anhydrous magnesium sulfate, then suspended

solids, and evaporated to dryness, afforded picric acid⁶ (5.15 g, 94%).

Isobutyryl Chloride (144)

Isobutyric acid (138.8 g, 1.15 molar) and thionyl chloride (243.1 g, 1.15 ml, 1.35 molar) were combined in a 300 ml round-bottomed flask equipped with a magnetic stirring bar and a reflux condenser. After the initial exothermic reaction had ceased, the mixture was gently refluxed for 5 hours. The reflux condenser was replaced by a fractionating column and the compound distilling at 80-82° was collected. Isobutyryl chloride (244.7 g, 80%) was obtained as a pungent, colorless liquid⁶ or (liquid film) 3000 cm^{-1} (C=O); n_D^{20} 1.37 (liquid), d_4^{20} 0.81, n_D^{25} 1.35 (liquid), d_4^{25} 0.79. n_D^{20} 1.40.

Isobutyrylbenzoate (145)

A solution consisting of sodium (34.8 g, 1.500 molar) isobutyrate (138.8 g, 1.50 molar), and 300 ml of chloroform was prepared in a 300 ml three-necked round-bottomed flask equipped with an addition funnel, mechanical stirrer, and an ice water cooling bath. Isobutyryl chloride (415.8 g, 1.501 molar) in 100 ml of chloroform was added dropwise to the above sodium solution, then stirred at ambient temperature for 12 hours. The chloroform mixture was washed with dilute hydrochloric acid, 5% sodium carbonate, and, finally, with water. The organic layer was dried over anhydrous magnesium sulfate then evaporated to dryness. Recrystallization of the residue from

⁶ Identification was based on spectroscopic comparison with an authentic sample.

1,3,5-Trisubstituted-2,4,6-Trisubstitutedbenzophenone-triazine (I.41)

A solution of dimethylketone-8-phenylindole (1.00 g, 31.4 mmol) in 500 ml of anhydrous ether was prepared in a 100 ml round-bottomed flask equipped with a magnetic stirring bar, and chilled to 0-5° with an ice water bath. This solution was treated with methyl fluoroacetate (1.62 ml, 2.31 g, 31.2 mmol), then stirred at ambient temperature for 24 hours. The solution mixture was decanted from the small amount of brown oil (0.50 g) adhering to the walls of the flask and evaporated at reduced pressure to a colorless crystalline solid (0.40 g). The crude triazine was recrystallized from hot absolute ethanol to yield 1,3,5-trisubstituted-2,4,6-trisubstitutedbenzophenone-triazine (0.30 g, 10%) as colorless prisms mp 149-151° (lit.¹³ mp 143-144°), λ_{max} (CHCl₃) 2815 m μ ¹⁴ (29-3-6), ν_{max} (CHCl₃) δ 1.62 (singlet, 18H), 6.32-7.12 (multiplet, 18H), δ_{H} (solid 1:1 C₁₂H₁₀N₃) 4.05 (4H), (liquid) 4.15 (4H).

Solid Catalyst Synthesis of Dimethylketone-8-Phenylindole (25) to 1,3,5-Trisubstituted-2,4,6-Trisubstitutedbenzophenone-triazine (I.41)

Dimethylketone-8-phenylindole (0.50 g, 3.4 mmol) and 25 ml of anhydrous ether were combined in a 50 ml round-bottomed flask and stirred magnetically until dissolved. A single drop of fluoroacetic acid was added to this solution and the resulting mixture stirred at ambient temperature for 18 hours. Evaporation of the ether at reduced pressure left a pale yellow solid residue. Recrystallization from absolute ethanol afforded colorless crystals (0.20 g, 40%, mp 144-145°). Spectroscopic properties with that of an authentic sample identified this material as 1,3,5-trisubstituted-2,4,6-trisubstitutedbenzophenone-triazine.

1.43 (singlet, 4H), 2.44 (singlet, 4H), 1.43-1.35 (broad, s-1 H), 3H,
 1 (2-dial doublet, 1.5H)

Synthesis of 1,3,5-Triphenyl-2-L-tryptophyl-4,6-dihydroxyindane-3,4,5,6-tetrahydro- α -trioxane-2-trioxane-1,2,3,4-tetraol (192)

1,3,5-Triphenyl-2-L-tryptophyl-4,6-dihydroxyindane-3,4,5,6-tetrahydro- α -trioxane-1,2,3,4-tetraol (191) (2.00 g, 3.12 mmol) and 50 ml of deionized water were stirred under reflux for a period of 11 hours. During this time, an insoluble white solid gradually appeared on the surface of the late yellow liquid phase. The white solid was collected on a small filter funnel and was shown by an analytical step 10 to be 1,3,5-triphenyl-2,4,6-trihydroxyindanepolyhydro- α -trioxane (192) (g. 192).

1,3,5-Triphenyl-2-L-tryptophyl-4,6-dihydroxyindane-3,4,5,6-tetrahydro- α -trioxane-1,2,3,4-tetraol (192)

1,3,5-Triphenyl-2-L-tryptophyl-4,6-dihydroxyindane-3,4,5,6-tetrahydro- α -trioxane-1,2,3,4-tetraol (1.45 g, 3.45 mmol), and 50 ml of ethanol were combined in a 125 ml volumetric flask equipped with a magnetic stirring bar. Addition of sodium hydroxide (0.210 g, 3.18 mmol) in several increments to the colored solution will initiate a violent reaction with attendant disappearance of the yellow indanone color component. After stirring for 15 minutes, 40 ml of water was added. 1,3,5-Triphenyl-2-L-tryptophyl-4,6-dihydroxyindane-3,4,5,6-tetrahydro- α -trioxane (1.14 g, 99%) was collected on a small filter funnel as a white powder. Recrystallization from hot benzene afforded white prisms which melted broadly at 150-161°; in (KBr) 1440 and 1270 cm^{-1} (C-O-H); and (KBr) δ 3.75 (aromatic,

distilled at 107°/0.25 mm Hg. 2-Methyl-5-*l*-prolylhistidinolyl chloride (1.36 g, 5.82 mmol) was obtained as a colorless liquid. IR (liquid film) 1690 cm^{-1} ($\text{N}=\text{O}$); ^1H NMR (CDCl_3) δ 1.44 (singlet, 3H, CH_3), 1.46 (d, 1H, CH_2), 1.56 (singlet, 3H, CH_3), 1.58 (singlet, 3H, CH_3), 2.34 (singlet, 3H, CH_3), 3.12-3.17 (multiplet, 4H).

The above prepared alkyl chloride (1.1 g, 37.5 mmol), 150 ml of toluene, and triethylamine (20.5 g, 91.5 ml, 0.375 mol) were added successively to a 100 ml round-bottomed flask equipped with a magnetic stirring bar, reflux condenser, and a nitrogen atmosphere. The mixture was stirred under reflux for 12 hours then filtered through a medium porosity filter to remove the precipitated triethylamine hydrochloride. The dark brown filtrate was concentrated at reduced pressure and vacuum distilled at 100-104°/0.45 mm Hg.

2-Methyl-5-*l*-prolylhistidine (1.36 g, 5.82 mmol) was obtained as a greenish yellow liquid. IR (liquid film) 1690 cm^{-1} ($\text{N}=\text{O}$); ^1H NMR (CDCl_3) δ 1.44 (singlet, 3H, CH_3), 1.46 (singlet, 3H, CH_3), 1.56 (singlet, 3H, CH_3), 1.58 (singlet, 3H, CH_3), 2.34 (singlet, 3H, CH_3), 3.12-3.17 (multiplet, 4H, CH_2). ^1H NMR (CDCl_3) 147.1 (NH), 147.1 (NH).

Methylation of 2-Methyl-5-*l*-Prolylhistidine (1.36 g, 5.82 mmol) with Methyl Chloroacetate. 1,3,5-Trib-4 α -methyl-2,4,6-tris-(2-methyl-5-*l*-prolylhistidinol)hydrochloride (1.36 g) and 1,3-Dimethyl-5-*l*-*l*-prolylhistidine-2-*l*-prolylhistidine-1,3,5-tris-(2-methyl-5-*l*-prolylhistidinol)hydrochloride (1.36 g).

A solution of 2-methyl-5-*l*-prolylhistidine (1.36 g, 5.82 mmol) in 50 ml of anhydrous ether was prepared in a 100 ml round-bottomed flask equipped with a magnetic stirring bar under a nitrogen atmosphere. Methyl chloroacetate (0.513 g, 5.82 ml, 0.46 mmol) was added, and the mixture stirred at room temperature for 12 hours. The ether was removed at reduced pressure and the pale yellow residue distilled with 20 ml of absolute ethanol. The colorless crystals of 1,3,5-trib-4 α -methyl-

sulfonic funnel, and a sodium sulfate drying tube, then added to 5-17 with an ion wire bath. 3,3-Diphenylacetyl chloride (82.0 g, 0.318 mole) in 75 ml of chloroform was added dropwise to the sodium solution and stirred at ambient temperature for 8 hours. The crude product mixture was washed with dilute hydrochloric acid, water, and, finally, dried over anhydrous magnesium sulfate. Evaporation of the solvent at reduced pressure afforded 8-isopropyl-3,3-diphenylacetamide (40.8 g, 48%) as a colorless solid. Recrystallization from hot benzene produced small colorless needles mp 133-135.5° (lit.¹⁰ mp 133-139°), n_D^{20} (CHCl₃) 1.450 (c=1.0) n_D^{25} (CHCl₃) 1.447 (doublet, J=7 Hz, 8H), 6.17 (septet, J=7 Hz, 1H), 5.35 (singlet, 1H), 5.45 (broad, 1H), 7.13 (doublet, 1H).

8-Isopropyl-3,3-diphenylchloride (XII)

8-Isopropyl-3,3-diphenylacetamide (40.8 g, 0.318 mole) and 50 ml of benzene were combined in a 100 ml round-bottomed flask equipped with a magnetic stirring bar and a reflux condenser. Phosphorus pentachloride (24.4 g, 75.0 mmol) was added to the white slurry, and the resulting suspension stirred under reflux for 30 minutes. The solvent and phosphoryl chloride were removed by evaporation at reduced pressure. Without further purification, the crude isobutyl chloride was transferred with 100 ml of toluene to a 100 ml round-bottomed flask equipped with a magnetic stirring bar, reflux condenser, and a sodium sulfate drying tube. The solution was treated with triethylamine (20.0 g, 0.488 mole, 15.5 ml), then stirred under reflux for 12 hours. Upon cooling, the mixture was poured through a column of activity 60/80 sand to remove the precipitated triethylamine hydrochloride. The filtrate was concentrated at reduced pressure to a brown liquid

8-(azepangyl-2-(2-methyl-6-azepangyl)oxane-3,3-diphenyl-4-benzophenyl)oxane-2-carboxylate (115)

8-(azepangyl-2-(2-methyl-6-azepangyl)oxane-3,3-diphenyl-4-benzophenyl)oxane-2-carboxylic acid (1.000 g, 2.103 mmol), and 30 ml of THF solvent were combined in a 100 ml volumetric flask and stirred magnetically. Triethylamine hydrochloride (0.128 g, 6.30 mmol) was added to the heterogeneous mixture in small increments over a period of 3 minutes. During this time, the dimethylformamide salt was rapidly consumed and was displaced by a fine milky white solid. After stirring at room temperature for an additional 15 minutes, the reaction was quenched with 50 ml of water. The solid was isolated by addition of hexane as a solid phase solvent, washed three times with 10 ml portions of water, and, finally, dried under vacuum. 8-(azepangyl-2-(2-methyl-6-azepangyl)oxane-3,3-diphenyl-4-benzophenyl)oxane-2-carboxylate (1.004 g, 74%) was obtained as a colorless powder. Recrystallization from petroleum ether (30-40°) afforded colorless prisms mp 119-120°, n_D^{20} (CHCl₃) 1.468 cm^{-1} (C=O-Ph), n_D^{20} (CHCl₃) 1.510 (broadest, 3-7 μ , 1760, 1.48 (broadest, 3-7 μ , 380, 1.52 (broadest, 30), 1.33-1.65 (multiplet, 20), 1.40 (broadest, 30), 1.30-1.65 (multiplet, 200), s/s (small for $\frac{C_{20}H_{24}N_2O_2}{446.5042}$) 446.5042.

Thermal data of 8-(azepangyl-2-(2-methyl-6-azepangyl)oxane-3,3-diphenyl-4-benzophenyl)oxane-2-carboxylate (115)

8-(azepangyl-2-(2-methyl-6-azepangyl)oxane-3,3-diphenyl-4-benzophenyl)oxane-2-carboxylate (0.10 g, 0.11 mmol), was packed into a low α , β , wax tube and measured in a system equipped with an inline T-tube kept cooled to -70° with a dry-ice/nitrogen bath. The sample containing the wax tube was heated with a heating burner flame until

Solvent	Amplification Factor (vol/vol)	
Pyridine	1:0	1:1
Hexylene chloride	1:1	1:0
Dichlorane	1:1	1:0
Benzene	1:0	1:0
Toluene	1:1	1:0

Preparation of 5-Isopropyl-2-(2-Methyl-5-Isopropylhexano-3,3-Diethyl-5-Hydroxyheptanoate) (III) in the Presence of Hexahydroxybenzotriazine

5-Isopropyl-2-(2-methyl-5-isopropylhexano-3,3-diethyl-5-hydroxyheptanoate) (I) (500 g., 0.005 mole) was packed into a 5 cm x 4 mm tube, then diluted with 0.5 ml of deuteriochloroform and dimethylsulfoxide-d₆ (5-10 ml., 0.005 g., 0.14 mole). The tube was capped, immersed in an oil bath preheated at 73°, and heated at this temperature for 30 minutes. The solution was filtered from the tube and evaporated at reduced pressure to remove the volatile components. The brown residue was taken up in deuteriochloroform and analyzed via NMR spectroscopy. Tetraphenylsilane was determined to be present in large amounts, but no 5-methyl signal was observed in the region δ 1.0-1.2 as expected for α 4H subject.

Preparation Hexahydroxybenzotriazine of 5-Isopropyl-2-(2-Methyl-5-Isopropylhexano-3,3-Diethyl-5-Hydroxyheptanoate) (III) from Dimethyl Sulfoxide

5-Isopropyl-2-(2-methyl-5-isopropylhexano-3,3-diethyl-5-hydroxyheptanoate) (I) (5.00 g., 0.14 mole) and 20 ml of dimethyl sulfoxide were placed in a 10 ml volumetric flask and heated to boiling. After a few minutes, the hot solution was filtered from the undissolved

action, and the clear solution obtained not suitable for crystallization. Several hours later, the fine colorless plates (1.18 g) were collected by vacuum filtration and dried in vacuo. Infrared (KBr disc) of this crystalline material and compared with reference spectra reported in the literature⁴⁸ identified the sample as tetraphenylsilane: mp 140-145.5° (dec.);⁴⁹ n_D^{20} 1.4933; n_D^{25} 1.4893; n_D^{30} 1.4861 and 1.4830 cm^{-1} (aromatic stretch and breathing modes); ν_{max} (KBr) 5 1.17-1.26 (μm - cm^{-1}).

3-Diphenylpropyl Chloride

3-Diphenylpropyl acid (14.3 g, 0.116 mole) and thionyl chloride (28.3 g, 15.3 ml, 0.395 mole) were combined in a 100 ml round-bottomed flask equipped with a reflux condenser and a magnetic stirring bar. After the initial exothermic reaction subsided, the mixture was refluxed for approximately 90 minutes. The reaction was quenched by a freezeflowing column and the compound distilling at 110-115° (lit.⁴⁹ bp 138-139°/130 mm Hg) was collected. 3-Diphenylpropyl chloride (14.7 g, 94%) was obtained as a pungent, colorless liquid. It distilled (bath 120-125°/150 mm Hg) and n_D^{20} 1.495 (refractive, 2-1 mm, 20°), 1.487 (refractive, 30°), 1.477 (refractive, 40°), 1.463 (refractive, 40°), 1.453 (refractive, 40°).

3-Diphenylpropyl-2-Hydroxybutyrate (114)

Diphenylacetic acid (19.46 g, 0.116 mole) and triethylamine (15.3 ml, 15.3 ml, 0.116 mole) were combined in a 100 ml round-bottomed flask equipped with an addition funnel and a magnetic stirring bar. 300 ml of chloroform was added and the mixture cooled to 0-5° with an ice

water bath. Triisopropylamide chloride (17.5 g, 0.086 mole) in 50 ml of chloroform was added dropwise to the stirred mixture, then stirred at ambient temperature for 4 hours. The mixture was concentrated to a 100 ml evaporatory residue and treated twice with dilute hydrochloric acid, then with water. The organic layer was dried over anhydrous magnesium sulfate and concentrated at reduced pressure to yield 2-isopropyl-2-ethylhexanamide (13.6 g, 68%) as a colorless waxy solid mp 14-16° (lit.)¹¹ mp 14-16°; ν (CHCl₃) 1650 cm⁻¹ (C=O), and (CHCl₃) + 2,2,4,4-tetrahydrofuran, 340, 1.44 (doublet, 2H, H_a, 4H), 1.25-1.35 (multiplet, 8H), 4.15 (quartet, 2-H, H_b, 3H), 5.55 (quintet, 4H).

Diisobutyl-2-isopropylamide, C₁₁H₂₃N

A solution of triphosphorophosgene (5.25 g, 0.05 mole) in 100 ml of methylene chloride was prepared in a 100 ml round-bottomed flask and chilled to 0-5° with an ice water bath. Triamine (5.00 g, 0.05 mole) was added dropwise to the stirred triphosphorophosgene solution until the first appearance of hydrogen color. At this point, diisopropyl-2-isopropylamide (5.00 g, 0.05 mole) and isobutylamine (5.00 g, 0.05 mole) were introduced to the mixture. The ice water bath was removed, replaced by a heating mantle, and the reaction flask fitted with a reflux condenser. The suspension was stirred and refluxed gently under a nitrogen atmosphere for a period of 40 minutes. During this time the mixture became homogeneous and assumed a dark brown appearance. After cooling to room temperature, the methylene chloride was evaporated at reduced pressure and the solid residue triturated twice with 10 ml portions of petroleum ether (20-40°). The pale yellow solids were washed and concentrated to a yellow liquid.

440, 1.31 (doublet, 2H), 1.4 (3.75 (multiplet, 2H), 2.22 (multiplet, 1H),
w/o (anal. for $C_{11}H_{12}N_2O_2$) 294.2834, (found) 294.2819)

3,5-Dimethylisobutyramide (11b)

Isobutyramide (11.3 g, 0.211 mole), triethylamine (29.3 g, 0.261 mole), and 150 ml of benzene were combined in a 500 ml three-necked round-bottomed flask equipped with a mechanical stirrer, and an ice water cooling bath. Triethylyl chloride (29.8 g, 0.188 mole) in 50 ml of benzene was added dropwise to the water solution. The mixture was stirred at ambient temperature for 4 hours, and then filtered from the triethylamine hydrochloride salt. The filtrate was concentrated at reduced pressure to a yellow liquid which on vacuum distillation at 126°/mm Hg 28 to 29 (lit.¹¹ 116-117°) afforded 3,5-dimethylisobutyramide (11.3 g, 80) as a colorless liquid.
1H (CDCl₃) 1440 cm⁻¹ (C=O), see (CDCl₃) 1.11 (doublet, 3-H, 80), 1.14 (multiplet, 3-H, 10), 1.17 (quint doublet, 1H,

3,5-Dimethyl-4-chloro-3-methyl-2-propionylamide (11c)

A solution of 3,5-dimethylisobutyramide (20.0 g, 0.201 mole), in 50 ml of anhydrous ether was prepared in a 500 ml three-necked round-bottomed flask equipped with a mechanical stirrer, addition funnel, and a nitrogen atmosphere. Propyl chloride (12.4g, 0.261 mole) in 25 ml of anhydrous ether was added dropwise to the water solution. During this time, rapid evolution of gases and the precipitation of a thick white solid occurred. The heterogeneous mixture was filtered with ether and stirred under nitrogen for approximately 21 hours. Triethylamine in 25 ml of anhydrous ether was added dropwise to the

white solid, and then stirred for an additional 12 hours. The unreacted product mixture was filtered from the precipitated sodium sulfate and concentrated at reduced pressure to a brown liquid. Vacuum distillation at 10⁻²/21 mm Hg (114-115) by 10⁻²/12 mm Hg produced 8,8-dimethyl-2-chloro-2-methyl-4-propenyltetrahydro-2H-pyran, 69% as a colorless, homogeneous liquid: d_4^{20} 1.05 cm³ 10-6-80, n_D^{20} 1.431, n_D^{25} 1.427 (singlet, 4.28 (singlet).

Preparation of 8,8-Dimethyl-2-chloro-2-methyl-4-propenyltetrahydro-2H-pyran (12.0)
Yield: 100% (100% Conversion)

A solution of sodium bis(trimethylsilyl)acetate (1.75 g, 15.8 mmol) in 50 ml of hexane was prepared in a 150-ml round-bottom flask equipped with an addition funnel, magnetic stirring bar, and a sodium sulfate drying tube. 8,8-Dimethyl-2-chloro-2-methyl-4-propenyltetrahydro-2H-pyran (2.00 g, 15.8 mmol) in 25 ml of hexane was added dropwise over a period of 5 minutes to the stirred allylamine solution. After stirring for 5 hours, the mixture was washed twice with 50 ml portions of water, dried over anhydrous magnesium sulfate, then evaporated to an oily residue (5.15 g). Vacuum distillation at 10⁻²/16.25 mm Hg produced 8,8-dimethyl-2-chloro-2-methyl-4-propenyltetrahydro-2-methyl-4-propenyltetrahydro-2H-pyran (2.47 g, 74%) as a colorless solid: mp 34-35°C; d_4^{20} 1.05 cm³ 10-6-80, n_D^{20} 1.431 (singlet, 1.05), 1.43 (doublet, 1.05), 1.43 (singlet, 1.05); n_D^{25} 1.427 (for C₁₂H₂₀ClO) 1.427-1.428, 1.427-1.428.

Treatment of 8,8-Dimethyl-1-1-Deuterio-1-Methyl-3-Propenylamine (III) with Nitrogen 1-Tetrazole

A suspension of potassium 1-tetrazole (5.10 g, 44.8 mmol), in 75 ml of dry benzene was prepared in a 150 ml round-bottomed flask equipped with a reflux condenser, magnetic stirring bar, and a nitrogen atmosphere. 8,8-Dimethyl-1-deutero-1-methyl-3-propenylamine (1.00 g, 12.4 mmol) in 10 ml of benzene was syringed into the tetrazolate suspension. This mixture was then refluxed under nitrogen for 12 hours. The product mixture was washed once with 50 ml of water, dried over anhydrous magnesium sulfate, and evaporated at reduced pressure to a pale yellow liquid. Vacuum distillation at 30-35°/0.15 mm Hg produced a colorless distillate (1.48 g) which contained pure 8,8-dimethyl-1-tetrazolide (ca. 10%) and a substance whose spectral properties are in agreement with 8,8-dimethyl-2-(1-tetrazol-2-ylmethyl)-3-propenylamine (λ_{max} 332), or (liquid film) 3000 cm^{-1} (C-C-H), and $\nu(\text{NCl}_2)$ = 1.34 (strong, 90), 1.33 (strong, 90), 1.12 (strong, 30), 1.01 (strong, 90), 901

Treatment of Dimethylamino-8,8-Dimethyl-1-tetrazolium Hexafluoroantimonate (III) with Nitrogen 1-Tetrazolylmethylamine

A solution of sodium dimethylallylamine (1.85 g, 11.7 mmol) in 11 ml of anhydrous ether was prepared in a 100 ml round-bottomed flask equipped with an addition funnel, magnetic stirring bar, and a nitrogen atmosphere. In a separate vessel, 8,8-dimethyl-1-deutero-1-methyl-3-propenylamine (1.00 g, 12.4 mmol) was added to a solution of silver perchlorate (1.10 g, 11.7 mmol) in 15 ml anhydrous ether. Precipitation of silver chloride occurred immediately. This heterogeneous mixture was rapidly transferred to the addition funnel

gig reaction filtration through a bed of Cellin. The clear dimethylbenzene-*N,N*-dimethylacetamide solution was added dropwise to the stirred ethylamide base solution. After 2 hours of stirring under nitrogen, the mixture was washed once with 50 ml of water, dried over anhydrous magnesium sulfate and concentrated in reduced pressure. Spectroscopic (NMR) analysis of the residue indicated that *N,N*-dimethylacetamide and *N,N*-dimethyl-1-*N,N*-bis(trimethylsilyl)-*N*-methyl-1-propenylamine were present in the approximate proportions of 1:1.1.

Preparation of Dimethylbenzene-*N,N*-Dimethylacetamide-Tetrahydrofuran (1:1) with Triethylamine

A suspension of phenol-*t*-butoxide (1.25 g, 11.2 mmol) in 25 ml of dry toluene was prepared in a 100 ml round-bottomed flask equipped with an addition funnel, magnetic stirring bar, and nitrogen atmosphere. In a separate vessel, *N,N*-dimethyl-1-chloro-2-methyl-1-propenylamine (1.58 g, 11.2 mmol) was added to a solution of silver tetrafluoroborate (2.10 g, 11.2 mmol) in 25 ml of toluene. Insoluble precipitation of silver chloride occurred. This heterogeneous mixture was rapidly transferred to the addition funnel gig reaction filtration through a bed of Cellin. The clear dimethylbenzene-*N,N*-dimethylacetamide-tetrahydrofuran solution was added dropwise to the stirred *t*-butoxide base slurry. After approximately 2 hours of stirring under nitrogen, the mixture was filtered, washed twice with 50 ml portions of water, dried over anhydrous magnesium sulfate, and concentrated in reduced pressure to a yellow residue. Spectroscopic (NMR) analysis of this residue indicated that *N,N*-dimethylacetamide and *N,N*-dimethyl-1-(2-methyl-1-propenyl)-1-*N,N*-bis(trimethylsilyl)-1-propenylamine were the only products.

8,8-Dimethyl-1,1-Dihydrocyclopent-1-ene

A solution consisting of triethylamine (13.8 g, 8.134 mol), dimethylamine (4.92 g, 8.134 mol), and 500 ml of chloroform was prepared in a 1000 ml round-bottomed flask equipped with an addition funnel, mechanical stirrer, and an ice-water cooling bath. Phosphoryl chloride (35.3 g, 4.134 mol) in 75 ml of chloroform was slowly added to the mixer slurry over a period of 15 minutes. The ice bath was removed, and the reaction mixture was stirred at ambient temperature for 2 hours. After quenching with dilute hydrochloric acid and water, the solution was dried over anhydrous organic solvents and then evaporated at reduced pressure to a solid residue. Recrystallization of this material from benzene-petroleum ether (40-120°) afforded 8,8-dimethyl-1,1-dihydrocyclopent-1-ene (30.2 g, 80%) as a colorless crystalline solid: mp 126-128° (lit. 89-90° (30-32°), in 4000₂) 1645 cm⁻¹ (C=O, var 1600₂), d 1.23 (singlet, 3H, 1.17 (singlet, 3H, 1.30 (singlet, 1H), 7.14 (singlet, 2H).

8,8-Dimethyl-1-Chloro-2,3-Dihydrocyclopent-1-ene

A solution of 8,8-dimethyl-1,1-dihydrocyclopent-1-ene (35.0 g, 41.4 mol) in 50 ml of chloroform was poured in a 500 ml round-bottomed flask equipped with a magnetic stirring bar and a reflux condenser. Phosphorus pentachloride (34.0 g, 41.4 mol) was added and the resulting slurry stirred under reflux for 1 hour. The solvent and phosphoryl chloride were removed at reduced pressure and the resulting mass of phosphoryl chloride chased with 50 ml of chloroform. The residue was treated with a mixture of triethylamine (6.26 g, 38.6 ml, 46.8 mol) in 50 ml of chloroform, after stirring

Trazen distillation at $122.5^{\circ}/75.30$ mm Hg produced 8-methyl-8-

isopropylisobutyramide (17.8 g, 82%) as a colorless liquid: d_4^{20} 1.0494 (lit.¹ 1.0510 cm^{-1} (9-10), n_D^{20} 1.435 (lit.¹ 1.435) (singlet, 2-7 Hz, 4H), 3.40 (singlet, 2-7 Hz, 2H), 3.15 (singlet, 2H), 4.58 (singlet, 2H), 7.3-7.4 (multiplet, 2H); n_D^{20} 1.4354 for $C_{12}H_{21}NO$ 180-1805, d_4^{20} 1.0510, 1.0505.

8-Methyl-8-Isopropyl-5-Oxobutyl-2-Methyl-2-Propyloxazolidin-2(1H)

A solution of 8-methyl-8-isopropylisobutyramide (28.7 g, 12.3 mmol) in 25 ml of anhydrous ether was prepared in a flask fitted 500 ml three-necked round-bottomed flask equipped with a mechanical stirrer, addition funnel, and a nitrogen atmosphere. Methyl chloran (0.44 g, 12.3 mmol) in 50 ml of anhydrous ether was added dropwise to the white solution then stirred for 14 hours. During this time, slow collection of gases took place with the aspiration of an oily liquid from the sidearm. Triethylamine (1.78 g, 12.3 mmol) in 25 ml of ether was added slowly to the above mixture and the suspension stirred for 1 hour before being filtered through a bed of Celite. The filter cake was washed with ether and the combined filtrates evaporated at reduced pressure to an white liquid. Trazen distillation at $72^{\circ}/8.5$ mm Hg produced 8-methyl-8-isopropyl-5-oxobutyl-2-methyl-2-propyloxazolidin-2-one (2.49 g, 44%) as a yellow liquid: d_4^{20} 1.0494 (lit.¹ 1.0494) cm^{-1} (9-10), n_D^{20} 1.435 (lit.¹ 1.435) (singlet, 2H), 3.38 (singlet, 2H), 3.78 (singlet, 2H), 7.3-7.4 (multiplet, 2H); n_D^{20} 1.4354 for $C_{12}H_{21}NO$ 180-1805, d_4^{20} 1.0510, 1.0505.

12-0₂, 1468 12-0-0₂; nmr (CDCl_3) δ 1.01 (triplet, 3_H, CH_3), 1.72 (singlet, 3_H), 1.84 (singlet, 3_H), 1.87 (singlet, 3_H), 1.97 (singlet, 3_H), 4.28 (quartet, 2_H, CH_2 , with CH_2CH_2 for C_2H_5 , CH_2CH_2 300, 2000, 1700) 281.0075.

A sample of the above α -chloroester was allowed to stand for 3 days in an open tube in solution with anhydrous diethylamine. Analysis of this solution via nmr spectroscopy showed the complete disappearance of signals associated with the α -chloroester. In addition no signals corresponding to 2-methyl-2-carboethoxymethyl-4-hydroxypentanoate, resulting from some hydrolysis, were apparent. There appeared lines at δ 1.47 (quartet, 2_H, CH_2), 1.52 (triplet, 3_H, CH_3), 2.88 (multiplet, 2_H, CH_2), 4.27 (singlet), 4.46 (quartet, 2_H, CH_2), and 5.15 (singlet). The structure has been assigned as 2-methyl-2-isopropyl-2-oxocyclopentanecarboxylate. Before hydrolysis reduction of an ethanolic solution of the above mixture produced, in addition to 2-methyl-2-carboethoxymethyl-4-hydroxypentanoate, a substance whose spectral properties are identical to those of ethyl 2-methyl-2-isobutyrylpentanoate.

Preparation of 2-Methyl-2-Carboethoxymethyl-2-Chloro-2-Methyl-2-Propionyl-2-Oxocyclopentanecarboxylate

A slurry of potassium trioxocarbonate (9.205 g., 4.45 mols) in 18 ml. of anhydrous ether was prepared in a 100 ml. round-bottomed flask equipped with an addition funnel, magnetic stirring bar, and a nitrogen atmosphere. A solution of 2-methyl-2-carboethoxymethyl-2-chloro-2-methyl-2-propionyl-2-oxocyclopentanecarboxylate (9.205 g., 4.45 mols) in 18 ml. of anhydrous ether was added dropwise to the stirred suspension. Small yellow fumes appeared on each drop made contact with the base, but this coloration quickly faded. After 12 hours of refluxing, the

pyrrolidine (15.6 g, 0.174 mol) was added to the acidic solution. The mixture was stirred under reflux for 90 minutes then overnight at room temperature. The orange solution was evaporated at reduced pressure to remove solvent and phosphoryl chloride, then treated with a solution of triethylamine (27.8 g, 0.175 mol) in 100 ml of toluene. This mixture was refluxed for 2 hours. The solution was filtered free of the precipitated hydrochloride salts and evaporated at reduced pressure to a brown liquid. Vacuum distillation at 25-30°/0 mm Hg, D_{20}^{25} 1.16, n_D^{25} 1.5192/0 mm Hg) afforded 2-isopropyl-3-ethoxycarbonyl-5,6-d g, 81% as a colorless liquid. d_4^{25} 1.021, n_D^{25} 1.458, n_D^{20} 1.457, n_D^{15} 1.456, n_D^{10} 1.455, n_D^5 1.454, n_D^0 1.453, d_4^{25} 1.021, d_4^{20} 1.020, d_4^{15} 1.019, d_4^{10} 1.018, d_4^5 1.017, d_4^0 1.016, d_4^{25} 1.015, d_4^{20} 1.014, d_4^{15} 1.013, d_4^{10} 1.012, d_4^5 1.011, d_4^0 1.010, d_4^{25} 1.009, d_4^{20} 1.008, d_4^{15} 1.007, d_4^{10} 1.006, d_4^5 1.005, d_4^0 1.004, d_4^{25} 1.003, d_4^{20} 1.002, d_4^{15} 1.001, d_4^{10} 1.000, d_4^5 0.999, d_4^0 0.998, d_4^{25} 0.997, d_4^{20} 0.996, d_4^{15} 0.995, d_4^{10} 0.994, d_4^5 0.993, d_4^0 0.992, d_4^{25} 0.991, d_4^{20} 0.990, d_4^{15} 0.989, d_4^{10} 0.988, d_4^5 0.987, d_4^0 0.986, d_4^{25} 0.985, d_4^{20} 0.984, d_4^{15} 0.983, d_4^{10} 0.982, d_4^5 0.981, d_4^0 0.980, d_4^{25} 0.979, d_4^{20} 0.978, d_4^{15} 0.977, d_4^{10} 0.976, d_4^5 0.975, d_4^0 0.974, d_4^{25} 0.973, d_4^{20} 0.972, d_4^{15} 0.971, d_4^{10} 0.970, d_4^5 0.969, d_4^0 0.968, d_4^{25} 0.967, d_4^{20} 0.966, d_4^{15} 0.965, d_4^{10} 0.964, d_4^5 0.963, d_4^0 0.962, d_4^{25} 0.961, d_4^{20} 0.960, d_4^{15} 0.959, d_4^{10} 0.958, d_4^5 0.957, d_4^0 0.956, d_4^{25} 0.955, d_4^{20} 0.954, d_4^{15} 0.953, d_4^{10} 0.952, d_4^5 0.951, d_4^0 0.950, d_4^{25} 0.949, d_4^{20} 0.948, d_4^{15} 0.947, d_4^{10} 0.946, d_4^5 0.945, d_4^0 0.944, d_4^{25} 0.943, d_4^{20} 0.942, d_4^{15} 0.941, d_4^{10} 0.940, d_4^5 0.939, d_4^0 0.938, d_4^{25} 0.937, d_4^{20} 0.936, d_4^{15} 0.935, d_4^{10} 0.934, d_4^5 0.933, d_4^0 0.932, d_4^{25} 0.931, d_4^{20} 0.930, d_4^{15} 0.929, d_4^{10} 0.928, d_4^5 0.927, d_4^0 0.926, d_4^{25} 0.925, d_4^{20} 0.924, d_4^{15} 0.923, d_4^{10} 0.922, d_4^5 0.921, d_4^0 0.920, d_4^{25} 0.919, d_4^{20} 0.918, d_4^{15} 0.917, d_4^{10} 0.916, d_4^5 0.915, d_4^0 0.914, d_4^{25} 0.913, d_4^{20} 0.912, d_4^{15} 0.911, d_4^{10} 0.910, d_4^5 0.909, d_4^0 0.908, d_4^{25} 0.907, d_4^{20} 0.906, d_4^{15} 0.905, d_4^{10} 0.904, d_4^5 0.903, d_4^0 0.902, d_4^{25} 0.901, d_4^{20} 0.900, d_4^{15} 0.899, d_4^{10} 0.898, d_4^5 0.897, d_4^0 0.896, d_4^{25} 0.895, d_4^{20} 0.894, d_4^{15} 0.893, d_4^{10} 0.892, d_4^5 0.891, d_4^0 0.890, d_4^{25} 0.889, d_4^{20} 0.888, d_4^{15} 0.887, d_4^{10} 0.886, d_4^5 0.885, d_4^0 0.884, d_4^{25} 0.883, d_4^{20} 0.882, d_4^{15} 0.881, d_4^{10} 0.880, d_4^5 0.879, d_4^0 0.878, d_4^{25} 0.877, d_4^{20} 0.876, d_4^{15} 0.875, d_4^{10} 0.874, d_4^5 0.873, d_4^0 0.872, d_4^{25} 0.871, d_4^{20} 0.870, d_4^{15} 0.869, d_4^{10} 0.868, d_4^5 0.867, d_4^0 0.866, d_4^{25} 0.865, d_4^{20} 0.864, d_4^{15} 0.863, d_4^{10} 0.862, d_4^5 0.861, d_4^0 0.860, d_4^{25} 0.859, d_4^{20} 0.858, d_4^{15} 0.857, d_4^{10} 0.856, d_4^5 0.855, d_4^0 0.854, d_4^{25} 0.853, d_4^{20} 0.852, d_4^{15} 0.851, d_4^{10} 0.850, d_4^5 0.849, d_4^0 0.848, d_4^{25} 0.847, d_4^{20} 0.846, d_4^{15} 0.845, d_4^{10} 0.844, d_4^5 0.843, d_4^0 0.842, d_4^{25} 0.841, d_4^{20} 0.840, d_4^{15} 0.839, d_4^{10} 0.838, d_4^5 0.837, d_4^0 0.836, d_4^{25} 0.835, d_4^{20} 0.834, d_4^{15} 0.833, d_4^{10} 0.832, d_4^5 0.831, d_4^0 0.830, d_4^{25} 0.829, d_4^{20} 0.828, d_4^{15} 0.827, d_4^{10} 0.826, d_4^5 0.825, d_4^0 0.824, d_4^{25} 0.823, d_4^{20} 0.822, d_4^{15} 0.821, d_4^{10} 0.820, d_4^5 0.819, d_4^0 0.818, d_4^{25} 0.817, d_4^{20} 0.816, d_4^{15} 0.815, d_4^{10} 0.814, d_4^5 0.813, d_4^0 0.812, d_4^{25} 0.811, d_4^{20} 0.810, d_4^{15} 0.809, d_4^{10} 0.808, d_4^5 0.807, d_4^0 0.806, d_4^{25} 0.805, d_4^{20} 0.804, d_4^{15} 0.803, d_4^{10} 0.802, d_4^5 0.801, d_4^0 0.800, d_4^{25} 0.799, d_4^{20} 0.798, d_4^{15} 0.797, d_4^{10} 0.796, d_4^5 0.795, d_4^0 0.794, d_4^{25} 0.793, d_4^{20} 0.792, d_4^{15} 0.791, d_4^{10} 0.790, d_4^5 0.789, d_4^0 0.788, d_4^{25} 0.787, d_4^{20} 0.786, d_4^{15} 0.785, d_4^{10} 0.784, d_4^5 0.783, d_4^0 0.782, d_4^{25} 0.781, d_4^{20} 0.780, d_4^{15} 0.779, d_4^{10} 0.778, d_4^5 0.777, d_4^0 0.776, d_4^{25} 0.775, d_4^{20} 0.774, d_4^{15} 0.773, d_4^{10} 0.772, d_4^5 0.771, d_4^0 0.770, d_4^{25} 0.769, d_4^{20} 0.768, d_4^{15} 0.767, d_4^{10} 0.766, d_4^5 0.765, d_4^0 0.764, d_4^{25} 0.763, d_4^{20} 0.762, d_4^{15} 0.761, d_4^{10} 0.760, d_4^5 0.759, d_4^0 0.758, d_4^{25} 0.757, d_4^{20} 0.756, d_4^{15} 0.755, d_4^{10} 0.754, d_4^5 0.753, d_4^0 0.752, d_4^{25} 0.751, d_4^{20} 0.750, d_4^{15} 0.749, d_4^{10} 0.748, d_4^5 0.747, d_4^0 0.746, d_4^{25} 0.745, d_4^{20} 0.744, d_4^{15} 0.743, d_4^{10} 0.742, d_4^5 0.741, d_4^0 0.740, d_4^{25} 0.739, d_4^{20} 0.738, d_4^{15} 0.737, d_4^{10} 0.736, d_4^5 0.735, d_4^0 0.734, d_4^{25} 0.733, d_4^{20} 0.732, d_4^{15} 0.731, d_4^{10} 0.730, d_4^5 0.729, d_4^0 0.728, d_4^{25} 0.727, d_4^{20} 0.726, d_4^{15} 0.725, d_4^{10} 0.724, d_4^5 0.723, d_4^0 0.722, d_4^{25} 0.721, d_4^{20} 0.720, d_4^{15} 0.719, d_4^{10} 0.718, d_4^5 0.717, d_4^0 0.716, d_4^{25} 0.715, d_4^{20} 0.714, d_4^{15} 0.713, d_4^{10} 0.712, d_4^5 0.711, d_4^0 0.710, d_4^{25} 0.709, d_4^{20} 0.708, d_4^{15} 0.707, d_4^{10} 0.706, d_4^5 0.705, d_4^0 0.704, d_4^{25} 0.703, d_4^{20} 0.702, d_4^{15} 0.701, d_4^{10} 0.700, d_4^5 0.699, d_4^0 0.698, d_4^{25} 0.697, d_4^{20} 0.696, d_4^{15} 0.695, d_4^{10} 0.694, d_4^5 0.693, d_4^0 0.692, d_4^{25} 0.691, d_4^{20} 0.690, d_4^{15} 0.689, d_4^{10} 0.688, d_4^5 0.687, d_4^0 0.686, d_4^{25} 0.685, d_4^{20} 0.684, d_4^{15} 0.683, d_4^{10} 0.682, d_4^5 0.681, d_4^0 0.680, d_4^{25} 0.679, d_4^{20} 0.678, d_4^{15} 0.677, d_4^{10} 0.676, d_4^5 0.675, d_4^0 0.674, d_4^{25} 0.673, d_4^{20} 0.672, d_4^{15} 0.671, d_4^{10} 0.670, d_4^5 0.669, d_4^0 0.668, d_4^{25} 0.667, d_4^{20} 0.666, d_4^{15} 0.665, d_4^{10} 0.664, d_4^5 0.663, d_4^0 0.662, d_4^{25} 0.661, d_4^{20} 0.660, d_4^{15} 0.659, d_4^{10} 0.658, d_4^5 0.657, d_4^0 0.656, d_4^{25} 0.655, d_4^{20} 0.654, d_4^{15} 0.653, d_4^{10} 0.652, d_4^5 0.651, d_4^0 0.650, d_4^{25} 0.649, d_4^{20} 0.648, d_4^{15} 0.647, d_4^{10} 0.646, d_4^5 0.645, d_4^0 0.644, d_4^{25} 0.643, d_4^{20} 0.642, d_4^{15} 0.641, d_4^{10} 0.640, d_4^5 0.639, d_4^0 0.638, d_4^{25} 0.637, d_4^{20} 0.636, d_4^{15} 0.635, d_4^{10} 0.634, d_4^5 0.633, d_4^0 0.632, d_4^{25} 0.631, d_4^{20} 0.630, d_4^{15} 0.629, d_4^{10} 0.628, d_4^5 0.627, d_4^0 0.626, d_4^{25} 0.625, d_4^{20} 0.624, d_4^{15} 0.623, d_4^{10} 0.622, d_4^5 0.621, d_4^0 0.620, d_4^{25} 0.619, d_4^{20} 0.618, d_4^{15} 0.617, d_4^{10} 0.616, d_4^5 0.615, d_4^0 0.614, d_4^{25} 0.613, d_4^{20} 0.612, d_4^{15} 0.611, d_4^{10} 0.610, d_4^5 0.609, d_4^0 0.608, d_4^{25} 0.607, d_4^{20} 0.606, d_4^{15} 0.605, d_4^{10} 0.604, d_4^5 0.603, d_4^0 0.602, d_4^{25} 0.601, d_4^{20} 0.600, d_4^{15} 0.599, d_4^{10} 0.598, d_4^5 0.597, d_4^0 0.596, d_4^{25} 0.595, d_4^{20} 0.594, d_4^{15} 0.593, d_4^{10} 0.592, d_4^5 0.591, d_4^0 0.590, d_4^{25} 0.589, d_4^{20} 0.588, d_4^{15} 0.587, d_4^{10} 0.586, d_4^5 0.585, d_4^0 0.584, d_4^{25} 0.583, d_4^{20} 0.582, d_4^{15} 0.581, d_4^{10} 0.580, d_4^5 0.579, d_4^0 0.578, d_4^{25} 0.577, d_4^{20} 0.576, d_4^{15} 0.575, d_4^{10} 0.574, d_4^5 0.573, d_4^0 0.572, d_4^{25} 0.571, d_4^{20} 0.570, d_4^{15} 0.569, d_4^{10} 0.568, d_4^5 0.567, d_4^0 0.566, d_4^{25} 0.565, d_4^{20} 0.564, d_4^{15} 0.563, d_4^{10} 0.562, d_4^5 0.561, d_4^0 0.560, d_4^{25} 0.559, d_4^{20} 0.558, d_4^{15} 0.557, d_4^{10} 0.556, d_4^5 0.555, d_4^0 0.554, d_4^{25} 0.553, d_4^{20} 0.552, d_4^{15} 0.551, d_4^{10} 0.550, d_4^5 0.549, d_4^0 0.548, d_4^{25} 0.547, d_4^{20} 0.546, d_4^{15} 0.545, d_4^{10} 0.544, d_4^5 0.543, d_4^0 0.542, d_4^{25} 0.541, d_4^{20} 0.540, d_4^{15} 0.539, d_4^{10} 0.538, d_4^5 0.537, d_4^0 0.536, d_4^{25} 0.535, d_4^{20} 0.534, d_4^{15} 0.533, d_4^{10} 0.532, d_4^5 0.531, d_4^0 0.530, d_4^{25} 0.529, d_4^{20} 0.528, d_4^{15} 0.527, d_4^{10} 0.526, d_4^5 0.525, d_4^0 0.524, d_4^{25} 0.523, d_4^{20} 0.522, d_4^{15} 0.521, d_4^{10} 0.520, d_4^5 0.519, d_4^0 0.518, d_4^{25} 0.517, d_4^{20} 0.516, d_4^{15} 0.515, d_4^{10} 0.514, d_4^5 0.513, d_4^0 0.512, d_4^{25} 0.511, d_4^{20} 0.510, d_4^{15} 0.509, d_4^{10} 0.508, d_4^5 0.507, d_4^0 0.506, d_4^{25} 0.505, d_4^{20} 0.504, d_4^{15} 0.503, d_4^{10} 0.502, d_4^5 0.501, d_4^0 0.500, d_4^{25} 0.499, d_4^{20} 0.498, d_4^{15} 0.497, d_4^{10} 0.496, d_4^5 0.495, d_4^0 0.494, d_4^{25} 0.493, d_4^{20} 0.492, d_4^{15} 0.491, d_4^{10} 0.490, d_4^5 0.489, d_4^0 0.488, d_4^{25} 0.487, d_4^{20} 0.486, d_4^{15} 0.485, d_4^{10} 0.484, d_4^5 0.483, d_4^0 0.482, d_4^{25} 0.481, d_4^{20} 0.480, d_4^{15} 0.479, d_4^{10} 0.478, d_4^5 0.477, d_4^0 0.476, d_4^{25} 0.475, d_4^{20} 0.474, d_4^{15} 0.473, d_4^{10} 0.472, d_4^5 0.471, d_4^0 0.470, d_4^{25} 0.469, d_4^{20} 0.468, d_4^{15} 0.467, d_4^{10} 0.466, d_4^5 0.465, d_4^0 0.464, d_4^{25} 0.463, d_4^{20} 0.462, d_4^{15} 0.461, d_4^{10} 0.460, d_4^5 0.459, d_4^0 0.458, d_4^{25} 0.457, d_4^{20} 0.456, d_4^{15} 0.455, d_4^{10} 0.454, d_4^5 0.453, d_4^0 0.452, d_4^{25} 0.451, d_4^{20} 0.450, d_4^{15} 0.449, d_4^{10} 0.448, d_4^5 0.447, d_4^0 0.446, d_4^{25} 0.445, d_4^{20} 0.444, d_4^{15} 0.443, d_4^{10} 0.442, d_4^5 0.441, d_4^0 0.440, d_4^{25} 0.439, d_4^{20} 0.438, d_4^{15} 0.437, d_4^{10} 0.436, d_4^5 0.435, d_4^0 0.434, d_4^{25} 0.433, d_4^{20} 0.432, d_4^{15} 0.431, d_4^{10} 0.430, d_4^5 0.429, d_4^0 0.428, d_4^{25} 0.427, d_4^{20} 0.426, d_4^{15} 0.425, d_4^{10} 0.424, d_4^5 0.423, d_4^0 0.422, d_4^{25} 0.421, d_4^{20} 0.420, d_4^{15} 0.419, d_4^{10} 0.418, d_4^5 0.417, d_4^0 0.416, d_4^{25} 0.415, d_4^{20} 0.414, d_4^{15} 0.413, d_4^{10} 0.412, d_4^5 0.411, d_4^0 0.410, d_4^{25} 0.409, d_4^{20} 0.408, d_4^{15} 0.407, d_4^{10} 0.406, d_4^5 0.405, d_4^0 0.404, d_4^{25} 0.403, d_4^{20} 0.402, d_4^{15} 0.401, d_4^{10} 0.400, d_4^5 0.399, d_4^0 0.398, d_4^{25} 0.397, d_4^{20} 0.396, d_4^{15} 0.395, d_4^{10} 0.394, d_4^5 0.393, d_4^0 0.392, d_4^{25} 0.391, d_4^{20} 0.390, d_4^{15} 0.389, d_4^{10} 0.388, d_4^5 0.387, d_4^0 0.386, d_4^{25} 0.385, d_4^{20} 0.384, d_4^{15} 0.383, d_4^{10} 0.382, d_4^5 0.381, d_4^0 0.380, d_4^{25} 0.379, d_4^{20} 0.378, d_4^{15} 0.377, d_4^{10} 0.376, d_4^5 0.375, d_4^0 0.374, d_4^{25} 0.373, d_4^{20} 0.372, d_4^{15} 0.371, d_4^{10} 0.370, d_4^5 0.369, d_4^0 0.368, d_4^{25} 0.367, d_4^{20} 0.366, d_4^{15} 0.365, d_4^{10} 0.364, d_4^5 0.363, d_4^0 0.362, d_4^{25} 0.361, d_4^{20} 0.360, d_4^{15} 0.359, d_4^{10} 0.358, d_4^5 0.357, d_4^0 0.356, d_4^{25} 0.355, d_4^{20} 0.354, d_4^{15} 0.353, d_4^{10} 0.352, d_4^5 0.351, d_4^0 0.350, d_4^{25} 0.349, d_4^{20} 0.348, d_4^{15} 0.347, d_4^{10} 0.346, d_4^5 0.345, d_4^0 0.344, $d_4^{$

4.38 (quartet, J=7 Hz, 1H), 4.81 (doublet, 2H).

Reduction of 8-benzyloxy-8-oxooctyl-5-oxooctanoate (1144) with sodium borohydride. (1145)-8-benzyloxy-8-oxooctyl-5-oxooctanoate (1145)

A solution of 8-benzyloxy-8-oxooctyl-5-oxooctanoate (1144) (1.40 g, 3.77 mmol) in 15 ml of ethanol was prepared in a 125 ml. three-necked flask and chilled to 0-5° with an ice-salt bath. Sodium borohydride (0.91 g, 3.77 mmol) was added in several small increments and then stirred at reflux temperature for 30 minutes. The solution was concentrated in a vacuum at reduced pressure. This residue was taken up in 25 ml of methylene chloride and dried over anhydrous magnesium sulfate. Evaporation of the solvent at reduced pressure produced ethyl 8-benzyloxy-8-oxooctanoate (0.40 g, 55%) as a colorless liquid. Its analytical sample was prepared by vacuum distillation (bath/boiler) at 50-100°/1 mm Hg. n_D^{20} (CHCl₃) 1.4366 (lit.¹ 1.4367), n_D^{25} (CHCl₃) 1.4344 (lit.¹ 1.4345), n_D^{20} (C₆H₆) 1.4311 (lit.¹ 1.4312), n_D^{25} (C₆H₆) 1.4288.

1-benzylsuccinic acid (11)

1-benzylsuccinic acid was prepared from 1-benzylsuccinimide according to the method described by Day and Gaudin¹⁰⁰ by 50-55°/1 mm Hg. Its lit.¹⁰⁰ n_D^{20} (CHCl₃) 1.4400 (lit.¹⁰⁰ 1.4401), n_D^{25} (CHCl₃) 1.4373 (lit.¹⁰⁰ 1.4374).

Ethyl 1,3-bis(4-oxo-4-oxobutyl)-3-oxobutanoate (1150)

Ethyl 1,3-bis(4-oxo-4-oxobutyl)-3-oxobutanoate was prepared by the ketone condensation of benzaldehyde with ethyl chloroacetate according to the method described by Day.⁷⁸ mp 32-33.5° (lit.⁷⁸

mg (2-45%), 14 (5mg) 1.780 cm⁻¹, and (221_g) δ 1.66 (quartet, 2-3 Hs, 180), 1.58 (quartet, 2-3 Hs, 18, 8₂), 1.17 (quartet, 2-3 Hs, 18, 8₁), 4.21 (quartet, 1-2 Hs, 180), 4.2-4.4 (multiplet, 180).

Thermolysis (isomerization) of 1,4,1-trimethyl-1,2-diphenyl-2-oxirane-carboxylic acid (1,1,2) in carbon tetrachloride

1,4,1-trimethyl-1,2-diphenyl-2-oxirane-carboxylic acid (1,1,2) 0.5117 mmol and approximately 1 ml of carbon tetrachloride were placed in a 5 mm i.d. test tube, and sealed with a test tube cap. The mixture was heated for 24 hours at 80° by suspending the tube inside a flask of refluxing benzene so that the tube completely surrounded the carbon tetrachloride-oxirane solution. The mixture (inner distribution) was analyzed approximately by careful integration of the observed oxirane ring proton signals, and was shown to be approximately 54/29/17 cis and 36/29/35 trans.

trans	1,1,2	δ_1	δ 1.17 (2,3 Hs)
		δ_2	1.66 (2,3 Hs)
cis	8 ₁	δ_1	1.46 (2,3 Hs)
		δ_2	1.66 (2,3 Hs)

Thermolysis of 1,4,1-trimethyl-1,2-diphenyl-2-oxirane-carboxylic acid (1,1,2) in 2-butylenediole (1,1,2)

1,4,1-trimethyl-1,2-diphenyl-2-oxirane-carboxylic acid (1,1,2) 1.644 mmol and 2-butylenediole (1,1,2) 1.778 g, 5.218 mmol were placed in a 5 mm i.d. test tube, capped, then heated in a steam bath until the oxirane had completely dissolved. The mixture was heated for 24 hours at 80° by suspending the tube inside a flask of refluxing benzene so that the tube surrounded the 2-butylenediole-oxirane solution. The

later's results were analyzed quantitatively by converting the net spectrum in the region 4.5-5.0 μ . No signals expected of a jet which were observed. Instead, identification of the emission took place in the cases of approximately 100 μ and 120 μ lines as determined by careful integrations of the signals associated with the calibration line systems.

1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,92,93,94,95,96,97,98,99,100,101,102,103,104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,196,197,198,199,200,201,202,203,204,205,206,207,208,209,210,211,212,213,214,215,216,217,218,219,220,221,222,223,224,225,226,227,228,229,230,231,232,233,234,235,236,237,238,239,240,241,242,243,244,245,246,247,248,249,250,251,252,253,254,255,256,257,258,259,260,261,262,263,264,265,266,267,268,269,270,271,272,273,274,275,276,277,278,279,280,281,282,283,284,285,286,287,288,289,290,291,292,293,294,295,296,297,298,299,300,301,302,303,304,305,306,307,308,309,310,311,312,313,314,315,316,317,318,319,320,321,322,323,324,325,326,327,328,329,330,331,332,333,334,335,336,337,338,339,340,341,342,343,344,345,346,347,348,349,350,351,352,353,354,355,356,357,358,359,360,361,362,363,364,365,366,367,368,369,370,371,372,373,374,375,376,377,378,379,380,381,382,383,384,385,386,387,388,389,390,391,392,393,394,395,396,397,398,399,400,401,402,403,404,405,406,407,408,409,410,411,412,413,414,415,416,417,418,419,420,421,422,423,424,425,426,427,428,429,430,431,432,433,434,435,436,437,438,439,440,441,442,443,444,445,446,447,448,449,450,451,452,453,454,455,456,457,458,459,460,461,462,463,464,465,466,467,468,469,470,471,472,473,474,475,476,477,478,479,480,481,482,483,484,485,486,487,488,489,490,491,492,493,494,495,496,497,498,499,500,501,502,503,504,505,506,507,508,509,510,511,512,513,514,515,516,517,518,519,520,521,522,523,524,525,526,527,528,529,530,531,532,533,534,535,536,537,538,539,540,541,542,543,544,545,546,547,548,549,550,551,552,553,554,555,556,557,558,559,560,561,562,563,564,565,566,567,568,569,570,571,572,573,574,575,576,577,578,579,580,581,582,583,584,585,586,587,588,589,590,591,592,593,594,595,596,597,598,599,600,601,602,603,604,605,606,607,608,609,610,611,612,613,614,615,616,617,618,619,620,621,622,623,624,625,626,627,628,629,630,631,632,633,634,635,636,637,638,639,640,641,642,643,644,645,646,647,648,649,650,651,652,653,654,655,656,657,658,659,660,661,662,663,664,665,666,667,668,669,670,671,672,673,674,675,676,677,678,679,680,681,682,683,684,685,686,687,688,689,690,691,692,693,694,695,696,697,698,699,700,701,702,703,704,705,706,707,708,709,710,711,712,713,714,715,716,717,718,719,720,721,722,723,724,725,726,727,728,729,730,731,732,733,734,735,736,737,738,739,740,741,742,743,744,745,746,747,748,749,750,751,752,753,754,755,756,757,758,759,760,761,762,763,764,765,766,767,768,769,770,771,772,773,774,775,776,777,778,779,780,781,782,783,784,785,786,787,788,789,790,791,792,793,794,795,796,797,798,799,800,801,802,803,804,805,806,807,808,809,810,811,812,813,814,815,816,817,818,819,820,821,822,823,824,825,826,827,828,829,830,831,832,833,834,835,836,837,838,839,840,841,842,843,844,845,846,847,848,849,850,851,852,853,854,855,856,857,858,859,860,861,862,863,864,865,866,867,868,869,870,871,872,873,874,875,876,877,878,879,880,881,882,883,884,885,886,887,888,889,890,891,892,893,894,895,896,897,898,899,900,901,902,903,904,905,906,907,908,909,910,911,912,913,914,915,916,917,918,919,920,921,922,923,924,925,926,927,928,929,930,931,932,933,934,935,936,937,938,939,940,941,942,943,944,945,946,947,948,949,950,951,952,953,954,955,956,957,958,959,960,961,962,963,964,965,966,967,968,969,970,971,972,973,974,975,976,977,978,979,980,981,982,983,984,985,986,987,988,989,990,991,992,993,994,995,996,997,998,999,1000

Styryl, trans-1,3-bis(phenyl-5-oxocyclopentadienyl) 15.88 g, 15.88 mmol, 1-benzyl-5-oxocyclopentadiene 41.66 g, 24.66 mmol and 24 ml of carbon tetrachloride were added to a glass tube. The mixture was polished in a tube furnace placed at 80° and allowed to react, unfiltered at this temperature for a period of 80 hours. The tube was opened and the contents transferred to a round-bottom flask. The volatile components were removed by evaporation at reduced pressure. The oily polymer residue was chromatographed on 20% deactivated alumina with cyclohexane as eluent. Phenyl-5(phenyl-5-oxocyclopentadienyl)cyclopentadiene-8'- α -butylidene (7.36 g, 9.15) was collected as a yellow solid and evaporated in vacuo to a yellow oil: λ_{max} (CCl₄) 3308 (6.6-6); 3241 cm⁻¹ (C=O, ester); n_D^{20} (CCl₄) = 1.31 (liquid), 1-7 Br, 20, 4.31 (singlet, 2H), 4.5-5.37 (multiplet, 2H), same reaction with 20% D₂O.

acid Catalyzed Hydrolysis of Phenyl-*O*-Phenyl-*N*-Carbonylmethyl-
carbamate-*N'*-*N*-Isopropylamide (127), 2-Phenyl-2-(2-Phenyl-*N*-Carbonylmethyl-
carbamate-*N'*-*N*-isopropylamide) (128)

A solution consisting of phospho (8-phosphorooxopentaphosphonic acid) = sodiumhexafluorophosphate (7.50 g, 0.11 mol) in 50 ml of various

was prepared in a 115 ml three-neck flask containing a magnetic stirring bar. Five drops of water were added to this solution followed by three drops of concentrated hydrochloric acid. At this point, the mixture immediately developed a pinkish brown color. The solution was heated slowly to reflux and water added dropwise until the turbidity persisted. Neutralization with 10% sodium carbonate produced a brilliant yellow solution. The organic soluble components were extracted with carbon tetrachloride, dried over anhydrous magnesium sulfate, and concentrated at reduced pressure to a yellow orange oil. The oil was applied to a silica gel column and eluted first with chloroform/methane tetrachloride (25:75), and then with chloroform. The second fraction (chloroform) contained 2-phospho-3-O-phospho-6-methylthymidylpyrimidin-5'-ribitylaminic acid (3.47 g, 44%) as a pale yellow oil. Purify was established by the observation of a single spot on TLC (1% NaCl in CHCl_3) R_F 0.08, 0.10, 0.12, 0.15, 0.18, 0.20, 0.22, 0.25, 0.28, 0.30, 0.32, 0.35, 0.38, 0.40, 0.42, 0.45, 0.48, 0.50, 0.52, 0.55, 0.58, 0.60, 0.62, 0.65, 0.68, 0.70, 0.72, 0.75, 0.78, 0.80, 0.82, 0.85, 0.88, 0.90, 0.92, 0.95, 0.98, 1.00, 1.02, 1.05, 1.08, 1.10, 1.12, 1.15, 1.18, 1.20, 1.22, 1.25, 1.28, 1.30, 1.32, 1.35, 1.38, 1.40, 1.42, 1.45, 1.48, 1.50, 1.52, 1.55, 1.58, 1.60, 1.62, 1.65, 1.68, 1.70, 1.72, 1.75, 1.78, 1.80, 1.82, 1.85, 1.88, 1.90, 1.92, 1.95, 1.98, 2.00, 2.02, 2.05, 2.08, 2.10, 2.12, 2.15, 2.18, 2.20, 2.22, 2.25, 2.28, 2.30, 2.32, 2.35, 2.38, 2.40, 2.42, 2.45, 2.48, 2.50, 2.52, 2.55, 2.58, 2.60, 2.62, 2.65, 2.68, 2.70, 2.72, 2.75, 2.78, 2.80, 2.82, 2.85, 2.88, 2.90, 2.92, 2.95, 2.98, 3.00, 3.02, 3.05, 3.08, 3.10, 3.12, 3.15, 3.18, 3.20, 3.22, 3.25, 3.28, 3.30, 3.32, 3.35, 3.38, 3.40, 3.42, 3.45, 3.48, 3.50, 3.52, 3.55, 3.58, 3.60, 3.62, 3.65, 3.68, 3.70, 3.72, 3.75, 3.78, 3.80, 3.82, 3.85, 3.88, 3.90, 3.92, 3.95, 3.98, 4.00, 4.02, 4.05, 4.08, 4.10, 4.12, 4.15, 4.18, 4.20, 4.22, 4.25, 4.28, 4.30, 4.32, 4.35, 4.38, 4.40, 4.42, 4.45, 4.48, 4.50, 4.52, 4.55, 4.58, 4.60, 4.62, 4.65, 4.68, 4.70, 4.72, 4.75, 4.78, 4.80, 4.82, 4.85, 4.88, 4.90, 4.92, 4.95, 4.98, 5.00, 5.02, 5.05, 5.08, 5.10, 5.12, 5.15, 5.18, 5.20, 5.22, 5.25, 5.28, 5.30, 5.32, 5.35, 5.38, 5.40, 5.42, 5.45, 5.48, 5.50, 5.52, 5.55, 5.58, 5.60, 5.62, 5.65, 5.68, 5.70, 5.72, 5.75, 5.78, 5.80, 5.82, 5.85, 5.88, 5.90, 5.92, 5.95, 5.98, 6.00, 6.02, 6.05, 6.08, 6.10, 6.12, 6.15, 6.18, 6.20, 6.22, 6.25, 6.28, 6.30, 6.32, 6.35, 6.38, 6.40, 6.42, 6.45, 6.48, 6.50, 6.52, 6.55, 6.58, 6.60, 6.62, 6.65, 6.68, 6.70, 6.72, 6.75, 6.78, 6.80, 6.82, 6.85, 6.88, 6.90, 6.92, 6.95, 6.98, 7.00, 7.02, 7.05, 7.08, 7.10, 7.12, 7.15, 7.18, 7.20, 7.22, 7.25, 7.28, 7.30, 7.32, 7.35, 7.38, 7.40, 7.42, 7.45, 7.48, 7.50, 7.52, 7.55, 7.58, 7.60, 7.62, 7.65, 7.68, 7.70, 7.72, 7.75, 7.78, 7.80, 7.82, 7.85, 7.88, 7.90, 7.92, 7.95, 7.98, 8.00, 8.02, 8.05, 8.08, 8.10, 8.12, 8.15, 8.18, 8.20, 8.22, 8.25, 8.28, 8.30, 8.32, 8.35, 8.38, 8.40, 8.42, 8.45, 8.48, 8.50, 8.52, 8.55, 8.58, 8.60, 8.62, 8.65, 8.68, 8.70, 8.72, 8.75, 8.78, 8.80, 8.82, 8.85, 8.88, 8.90, 8.92, 8.95, 8.98, 9.00, 9.02, 9.05, 9.08, 9.10, 9.12, 9.15, 9.18, 9.20, 9.22, 9.25, 9.28, 9.30, 9.32, 9.35, 9.38, 9.40, 9.42, 9.45, 9.48, 9.50, 9.52, 9.55, 9.58, 9.60, 9.62, 9.65, 9.68, 9.70, 9.72, 9.75, 9.78, 9.80, 9.82, 9.85, 9.88, 9.90, 9.92, 9.95, 9.98, 10.00, 10.02, 10.05, 10.08, 10.10, 10.12, 10.15, 10.18, 10.20, 10.22, 10.25, 10.28, 10.30, 10.32, 10.35, 10.38, 10.40, 10.42, 10.45, 10.48, 10.50, 10.52, 10.55, 10.58, 10.60, 10.62, 10.65, 10.68, 10.70, 10.72, 10.75, 10.78, 10.80, 10.82, 10.85, 10.88, 10.90, 10.92, 10.95, 10.98, 11.00, 11.02, 11.05, 11.08, 11.10, 11.12, 11.15, 11.18, 11.20, 11.22, 11.25, 11.28, 11.30, 11.32, 11.35, 11.38, 11.40, 11.42, 11.45, 11.48, 11.50, 11.52, 11.55, 11.58, 11.60, 11.62, 11.65, 11.68, 11.70, 11.72, 11.75, 11.78, 11.80, 11.82, 11.85, 11.88, 11.90, 11.92, 11.95, 11.98, 12.00, 12.02, 12.05, 12.08, 12.10, 12.12, 12.15, 12.18, 12.20, 12.22, 12.25, 12.28, 12.30, 12.32, 12.35, 12.38, 12.40, 12.42, 12.45, 12.48, 12.50, 12.52, 12.55, 12.58, 12.60, 12.62, 12.65, 12.68, 12.70, 12.72, 12.75, 12.78, 12.80, 12.82, 12.85, 12.88, 12.90, 12.92, 12.95, 12.98, 13.00, 13.02, 13.05, 13.08, 13.10, 13.12, 13.15, 13.18, 13.20, 13.22, 13.25, 13.28, 13.30, 13.32, 13.35, 13.38, 13.40, 13.42, 13.45, 13.48, 13.50, 13.52, 13.55, 13.58, 13.60, 13.62, 13.65, 13.68, 13.70, 13.72, 13.75, 13.78, 13.80, 13.82, 13.85, 13.88, 13.90, 13.92, 13.95, 13.98, 14.00, 14.02, 14.05, 14.08, 14.10, 14.12, 14.15, 14.18, 14.20, 14.22, 14.25, 14.28, 14.30, 14.32, 14.35, 14.38, 14.40, 14.42, 14.45, 14.48, 14.50, 14.52, 14.55, 14.58, 14.60, 14.62, 14.65, 14.68, 14.70, 14.72, 14.75, 14.78, 14.80, 14.82, 14.85, 14.88, 14.90, 14.92, 14.95, 14.98, 15.00, 15.02, 15.05, 15.08, 15.10, 15.12, 15.15, 15.18, 15.20, 15.22, 15.25, 15.28, 15.30, 15.32, 15.35, 15.38, 15.40, 15.42, 15.45, 15.48, 15.50, 15.52, 15.55, 15.58, 15.60, 15.62, 15.65, 15.68, 15.70, 15.72, 15.75, 15.78, 15.80, 15.82, 15.85, 15.88, 15.90, 15.92, 15.95, 15.98, 16.00, 16.02, 16.05, 16.08, 16.10, 16.12, 16.15, 16.18, 16.20, 16.22, 16.25, 16.28, 16.30, 16.32, 16.35, 16.38, 16.40, 16.42, 16.45, 16.48, 16.50, 16.52, 16.55, 16.58, 16.60, 16.62, 16.65, 16.68, 16.70, 16.72, 16.75, 16.78, 16.80, 16.82, 16.85, 16.88, 16.90, 16.92, 16.95, 16.98, 17.00, 17.02, 17.05, 17.08, 17.10, 17.12, 17.15, 17.18, 17.20, 17.22, 17.25, 17.28, 17.30, 17.32, 17.35, 17.38, 17.40, 17.42, 17.45, 17.48, 17.50, 17.52, 17.55, 17.58, 17.60, 17.62, 17.65, 17.68, 17.70, 17.72, 17.75, 17.78, 17.80, 17.82, 17.85, 17.88, 17.90, 17.92, 17.95, 17.98, 18.00, 18.02, 18.05, 18.08, 18.10, 18.12, 18.15, 18.18, 18.20, 18.22, 18.25, 18.28, 18.30, 18.32, 18.35, 18.38, 18.40, 18.42, 18.45, 18.48, 18.50, 18.52, 18.55, 18.58, 18.60, 18.62, 18.65, 18.68, 18.70, 18.72, 18.75, 18.78, 18.80, 18.82, 18.85, 18.88, 18.90, 18.92, 18.95, 18.98, 19.00, 19.02, 19.05, 19.08, 19.10, 19.12, 19.15, 19.18, 19.20, 19.22, 19.25, 19.28, 19.30, 19.32, 19.35, 19.38, 19.40, 19.42, 19.45, 19.48, 19.50, 19.52, 19.55, 19.58, 19.60, 19.62, 19.65, 19.68, 19.70, 19.72, 19.75, 19.78, 19.80, 19.82, 19.85, 19.88, 19.90, 19.92, 19.95, 19.98, 20.00, 20.02, 20.05, 20.08, 20.10, 20.12, 20.15, 20.18, 20.20, 20.22, 20.25, 20.28, 20.30, 20.32, 20.35, 20.38, 20.40, 20.42, 20.45, 20.48, 20.50, 20.52, 20.55, 20.58, 20.60, 20.62, 20.65, 20.68, 20.70, 20.72, 20.75, 20.78, 20.80, 20.82, 20.85, 20.88, 20.90, 20.92, 20.95, 20.98, 21.00, 21.02, 21.05, 21.08, 21.10, 21.12, 21.15, 21.18, 21.20, 21.22, 21.25, 21.28, 21.30, 21.32, 21.35, 21.38, 21.40, 21.42, 21.45, 21.48, 21.50, 21.52, 21.55, 21.58, 21.60, 21.62, 21.65, 21.68, 21.70, 21.72, 21.75, 21.78, 21.80, 21.82, 21.85, 21.88, 21.90, 21.92, 21.95, 21.98, 22.00, 22.02, 22.05, 22.08, 22.10, 22.12, 22.15, 22.18, 22.20, 22.22, 22.25, 22.28, 22.30, 22.32, 22.35, 22.38, 22.40, 22.42, 22.45, 22.48, 22.50, 22.52, 22.55, 22.58, 22.60, 22.62, 22.65, 22.68, 22.70, 22.72, 22.75, 22.78, 22.80, 22.82, 22.85, 22.88, 22.90, 22.92, 22.95, 22.98, 23.00, 23.02, 23.05, 23.08, 23.10, 23.12, 23.15, 23.18, 23.20, 23.22, 23.25, 23.28, 23.30, 23.32, 23.35, 23.38, 23.40, 23.42, 23.45, 23.48, 23.50, 23.52, 23.55, 23.58, 23.60, 23.62, 23.65, 23.68, 23.70, 23.72, 23.75, 23.78, 23.80, 23.82, 23.85, 23.88, 23.90, 23.92, 23.95, 23.98, 24.00, 24.02, 24.05, 24.08, 24.10, 24.12, 24.15, 24.18, 24.20, 24.22, 24.25, 24.28, 24.30, 24.32, 24.35, 24.38, 24.40, 24.42, 24.45, 24.48, 24.50, 24.52, 24.55, 24.58, 24.60, 24.62, 24.65, 24.68, 24.70, 24.72, 24.75, 24.78, 24.80, 24.82, 24.85, 24.88, 24.90, 24.92, 24.95, 24.98, 25.00, 25.02, 25.05, 25.08, 25.10, 25.12, 25.15, 25.18, 25.20, 25.22, 25.25, 25.28, 25.30, 25.32, 25.35, 25.38, 25.40, 25.42, 25.45, 25.48, 25.50, 25.52, 25.55, 25.58, 25.60, 25.62, 25.65, 25.68, 25.70, 25.72, 25.75, 25.78, 25.80, 25.82, 25.85, 25.88, 25.90, 25.92, 25.95, 25.98, 26.00, 26.02, 26.05, 26.08, 26.10, 26.12, 26.15, 26.18, 26.20, 26.22, 26.25, 26.28, 26.30, 26.32, 26.35, 26.38, 26.40, 26.42, 26.45, 26.48, 26.50, 26.52, 26.55, 26.58, 26.60, 26.62, 26.65, 26.68, 26.70, 26.72, 26.75, 26.78, 26.80, 26.82, 26.85, 26.88, 26.90, 26.92, 26.95, 26.98, 27.00, 27.02, 27.05, 27.08, 27.10, 27.12, 27.15, 27.18, 27.20, 27.22, 27.25, 27.28, 27.30, 27.32, 27.35, 27.38, 27.40, 27.42, 27.45, 27.48, 27.50, 27.52, 27.55, 27.58, 27.60, 27.62, 27.65, 27.68, 27.70, 27.72, 27.75, 27.78, 27.80, 27.82, 27.85, 27.88, 27.90, 27.92, 27.95, 27.98, 28.00, 28.02, 28.05, 28.08, 28.10, 28.12, 28.15, 28.18, 28.20, 28.22, 28.25, 28.28, 28.30, 28.32, 28.35, 28.38, 28.40, 28.42, 28.45, 28.48, 28.50, 28.52, 28.55, 28.58, 28.60, 28.62, 28.65, 28.68, 28.70, 28.72, 28.75, 28.78, 28.80, 28.82, 28.85, 28.88, 28.90, 28.92, 28.95, 28.98, 29.00, 29.02, 29.05, 29.08, 29.10, 29.12, 29.15, 29.18, 29.20, 29.22, 29.25, 29.28, 29.30, 29.32, 29.35, 29.38, 29.40, 29.42, 29.45, 29.48, 29.50, 29.52, 29.55, 29.58, 29.60, 29.62, 29.65, 29.68, 29.70, 29.72, 29.75, 29.78, 29.80, 29.82, 29.85, 29.88, 29.90, 29.92, 29.95, 29.98, 30.00, 30.02, 30.05, 30.08, 30.10, 30.12, 30.15, 30.18, 30.20, 30.22, 30.25, 30.28, 30.30, 30.32, 30.35, 30.38, 30.40, 30.42, 30.45, 30.48, 30.50, 30.52, 30.55, 30.58, 30.60, 30.62, 30.65, 30.68, 30.70, 30.72, 30.75, 30.78, 30.80, 30.82, 30.85, 30.88, 30.90, 30.92, 30.95, 30.98, 31.00, 31.02, 31.05, 31.08, 31.10, 31.12, 31.15, 31.18, 31.20, 31.22, 31.25, 31.28, 31.30, 31.32, 31.35, 31.38, 31.40, 31.42, 31.45, 31.48, 31.50, 31.52, 31.55, 31.58, 31.60, 31.62, 31.65, 31.68, 31.70, 31.72, 31.75, 31.78, 31.80, 31.82, 31.85, 31.88, 31.90, 31.92, 31.95, 31.98, 32.00, 32.02, 32.05, 32.08, 32.10, 32.12, 32.15, 32.18, 32.20, 32.22, 32.25, 32.28, 32.30, 32.32, 32.35, 32.38, 32.40, 32.42, 32.45, 32.48, 32.50, 32.52, 32.55, 32.58, 32.60, 32.62, 32.65, 32.68, 32.70, 32.72, 32.75, 32.78, 32.80, 32.82, 32.85, 32.88, 32.90, 32.92, 32.95, 32.98, 33.00, 33.02, 33.05, 33.08, 33.10, 33.12, 33.15, 33.18, 33.20, 33.22, 33.25, 33.28, 33.30, 33.32, 33.35, 33.38, 33.40, 33.42, 33.45, 33.48, 33.50, 33.52, 33.55, 33.58, 33.60, 33.62, 33.65, 33.68, 33.70, 33.72, 33.75, 33.78, 33.80, 33.82, 33.85, 33.88, 33.90, 33.92, 33.95, 33.98, 34.00, 34.02, 34.05, 34.08, 34.10, 34.12, 34.15, 34.18, 34.20, 34.22, 34.25, 34.28, 34.30, 34.32, 34.35, 34.38, 34.40, 34.42, 34.45, 34.48, 34.50, 34.52, 34.55, 34.58, 34.60, 34.62, 34.65, 34.68, 34.70, 34.72, 34.75, 34.78, 34.80, 34.82, 34.85, 34.88, 34.90, 34.92, 34.95, 34.98, 35.00, 35.02, 35.05, 35.08, 35.10, 35.12, 35.15, 35.18, 35.20, 35.22, 35.25, 35.28, 35.30, 35.32, 35.35, 35.38, 35.40, 35.42, 35.45, 35.48, 35.50, 35.52, 35.55, 35.58, 35.60, 35.62, 35.65, 35.68, 35.70, 35.72, 35.75, 35.78, 35.80, 35.82, 35.85, 35.88, 35.90, 35.92, 35.95, 35.98, 36.00, 36.02, 36.05, 36.08, 36.10, 36.12, 36.15, 36.18, 36.20, 36.22, 36.25, 36.28, 36.30, 36.32, 36.35, 36.38, 36.40, 36.42, 36.45, 36.48, 36.50, 36.52, 36.55, 36.58, 36.60, 36.62, 36.65, 36.68, 36.70, 36.72, 36.75, 36.78, 36.80, 36.82, 36.85, 36.88, 36.90, 36.92, 36.95, 36.98, 37.00, 37.02, 37.05, 37.08, 37.10, 37.12, 37.15, 37.18, 37.20, 37.22, 37.25, 37.28, 37.30, 37.32, 37.35, 37.38, 37.40, 37.42, 37.45, 37.48, 37.50, 37.52, 37.55, 37.58, 37.60, 37.62, 37.65, 37.68, 37.70, 37.72, 37.75, 37.78, 37.80, 37.82, 37.85, 37.88, 37.90, 37.92, 37.95, 37.98, 38.00, 38.02, 38.05, 38.08, 38.10, 38.12, 38.15, 38.18, 38.20, 38.22, 38.25, 38.28, 38.30, 38.32, 38.35, 38.38, 38.40, 38.42, 38.45, 38.48, 38.50, 38.52, 38.55, 38.58, 38.60, 38.62, 38.65, 38.68, 38.70, 38.72, 38.75, 38.78, 38.80, 38.82, 38.85, 38.88, 38.90, 38.92, 38.95, 38.98, 39.00, 39.02, 39.05, 39.08, 39.10, 39.12, 39.15, 39.18, 39.20, 39.22, 39.25, 39.28, 39.30, 39.32, 39.35, 39.38, 39.40, 39.42, 39.45, 39.48, 39.50, 39.52, 39.55, 39.58, 39.60, 39.62, 39.65, 39.68, 39.70, 39.72, 39.75, 39.78, 39.80, 39.82, 39.85, 39.88, 39.90, 39.92, 39.95, 39.98, 40.00, 40.02, 40.05, 40.08, 40.10, 40.12, 40.15, 40.18, 40.20, 40.22, 40.25, 40.28, 40.30, 40.32, 40.35, 40.38, 40.40, 40.42, 40.45, 40.48, 40.50, 40.52, 40.55, 40.58, 40.60, 40.62, 40.65, 40.68, 40.70, 40.72, 40.75, 40.78, 40.80, 40.82, 40.85, 40.88, 40.90, 40.92, 40.95, 40.98, 41.00, 41.02, 41.05, 41.08, 41.10, 41.12, 41.15, 41.18, 41.20, 41.22, 41.25, 41.28, 41.30, 41.32, 41.35, 41.38, 41.40, 41.42, 41.45, 41.48, 41.50, 41.52, 41.55, 41.58, 41.60, 41.62, 41.65, 41.68, 41.70, 41.72, 41.75, 41.78, 41.80, 41.82, 41.85, 41.88, 41.90, 41.92, 41.95, 41.98, 42.00, 42.02, 42.05, 42.08, 42.10, 42.12, 42.15, 42.18, 42.20, 42.22, 42.25, 42.28, 42.30, 42.32, 42.35, 42.38, 42.40, 42.42, 42.45, 42.48, 42.50, 42.52, 42.55, 42.58, 42.60, 42.62, 42.65, 42.68, 42.70, 42.72, 42.75, 42.78, 42.80, 42.82, 42.85, 42.88, 42.90, 42.92, 42.95, 42.98, 43.00, 43.02, 43.05, 43.08, 43.10, 43.12, 43.15, 43.18, 43.20, 43.22, 43.25, 43.28, 43.30, 43.32, 43.35, 43.38, 43.40, 43.42, 43.45, 43.48, 43.50, 43.52, 43.55, 43.58, 43.60, 43.62, 43.65, 43.68, 43.70, 43.72, 43.75, 43.78, 43.80, 43.82, 43.85, 43.88, 43.90, 43.92, 43.95, 43.98, 44.00, 44.02, 44.05, 44.08, 44.10, 44.12, 44.15, 44.18, 44.20, 44.22, 44.25, 44.28, 44.30, 44.32, 44.35, 44.38, 44.40, 44.42, 44.45, 44.48, 44.50, 44.52, 44.55,

1380 cm^{-1} ($\nu_{\text{C=O}}$, carb. carbonyl) + 3.30 (singlet, 2H), + $4.5\text{--}7.30$ (multiplet, 4H).

2-bromo-2-(phenylacetyl) bromide (242)

Phenylacetyl chloride (14.3 g , 51.8 mmol) and bromine (23.9 g , 104 mmol) were combined in a 100 ml round-bottomed flask equipped with a reflux condenser and a small magnetic stirring bar. The flask was immersed in a silicon oil bath pre-set at 130° , and the mixture stirred at this temperature for approximately 50 minutes. The product mixture was vacuum distilled at $30^\circ/\text{mm}$, 0.45 mm Hg ($140\text{--}150^\circ/10\text{ mm Hg}$) to give 2-bromo-2-(phenylacetyl) bromide (11.8 g , 33%) as a colorless liquid. IR (liquid film) 1700 cm^{-1} ($\nu_{\text{C=O}}$, carb. carbonyl) + 3.17 (singlet, 0.5H), 3.40 (singlet, 0.5H), + $7.15\text{--}7.3$ (multiplet, 4H).

2-bromo-2-(phenylisobutyryl) bromide (243)

A solution consisting of isobutyranol (3.81 g , 8.46 mmol), isobutyranol (9.08 g , 11.1 mmol , 79.1 mmol) and 230 ml of benzene was prepared in a 100 ml three-necked round-bottomed flask equipped with an addition funnel and a mechanical stirrer. This mixture was stirred and cooled to $5\text{--}5^\circ$ with an ice water bath while a solution of 2-bromo-2-(phenylacetyl) bromide (23.6 g , 51.8 mmol) in 75 ml of benzene was added dropwise over a period of 30 minutes. After stirring for an additional 30 minutes, the isobutyranol isobutyranol was removed by vacuum filtration. The filtrate was washed with dilute hydrochloric acid, dried over anhydrous magnesium sulfate, concentrated at reduced pressure, and left under for crystallization. 2-bromo-2-(phenyl-2-isobutyryl) bromide (16.7 g , 34%) was obtained as small colorless prisms.

Hydrochloride (see scheme provided as original sample) mp 107-108°C; IR (KBr) 3315 (OH, stretch), 1628 cm^{-1} (C=O); ^1H NMR (CDCl_3) δ 1.18 (triplet, 3H, 3-CH₃), 2.31 (singlet, 3H, 4-CH₃), 3.42 (quartet, 2H, 2-CH₂), 4.42 (quartet, 2H, 1-CH₂), 7.18-7.58 (multiplet, 5H).

ANAL. Calcd for $\text{C}_{12}\text{H}_{15}\text{NO}_2$: C, 71.15; H, 7.51; N, 1.18. Found: C, 71.45; H, 7.89; N, 1.46. IR, 1612 cm^{-1} .

Ethyl 8-Phenylglycidate (14.5)

Ethyl bromoacetate (15.3 g, 8.165 mol), sodium (23.8 g, 1.04 mol) and 180 ml of benzene were combined in a 250 ml round-bottom flask equipped with a reflux condenser and a magnetic stirring bar. The mixture was stirred under reflux for 11 hours, cooled to room temperature, then filtered from the precipitated sodium hydroxide (15.3 g, 8.165 mol, 96.4%). The filtrate was evaporated to dryness, and the residue dissolved in a minimum amount of diethyl ether. Ethyl 8-phenylglycidate (14.5 g, 82%) was produced as a pale yellow crystalline solid; mp 33-34°C (lit.¹⁸ mp 32°C); IR (CDCl_3) 3315 (OH, stretch), 1733 cm^{-1} (C=O); ^1H NMR (CDCl_3) δ 1.12 (triplet, 3H, 3-CH₃), 1.75 (triplet, 3H, 4-CH₃), 2.41 (quartet, 2H, 2-CH₂), 4.24-4.32 (multiplet, 2H).

Independent Synthesis of 8-Phenyl-2-[5-Phenyl-2-Carboxyhexanoate]ethyl 8-Phenylglycidate

Ethyl 8-phenylglycidate (5.00 g, 25.0 mmol), 5-bromo-2-phenyl-2-oxopentanoate (4.30 g, 16.7 mmol), triethylamine (1.79 g, 16.7 mmol), and 15 ml of benzene were combined in a 250 ml round-bottom flask and refluxed for four days. The precipitated triethylamine hydrobromide was filtered by vacuum filtration. The filtrate was washed with dilute aqueous sodium carbonate, and, finally, dried over

anhydrous magnesium sulfate). Recrystallization of the solvent at reduced pressure afforded 1-phenyl-1-(2-phenyl-6-methylheptyl)indole-3'-
 1-ynylacetamide (0.80 g, 98%) as a pale yellow oil. An analytically pure sample was prepared by chromatographing approximately 1.0 g of the above solids on a silica gel column packed in chloroform-carbon tetrachloride (15:25). Nine 20-ml. fractions were collected by eluting with chloroform-carbon tetrachloride (15:25), the last three of which were shown by TLC analysis to contain products completely free of unreacted starting material. These were combined, evaporated at reduced pressure, and their combined oil high vacuum. The resulting yellow oil exhibited the following properties: IR (CHCl_3) 3075 (br, aromatic), 2120 ($\text{C}\equiv\text{C}$, weak), 1670 ($\text{C}=\text{O}$, weak), 1540 (cm^{-1} (sh), broad, str. COC_2H_5) & 1.40 (triplet, 3H, t, 3H), 1.10 (singlet, 3H, s, 3H) (singlet, 3H), 4.50 (quartet, 3H, t, 3H), 1.10 (singlet, 3H, s, 3H) (singlet, 3H), 4.30-7.30 (multiplet, 10H), 8.10 (broad, 3H).

ANAL. Calcd for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}$: C, 79.45; H, 7.00; N, 3.55.

Found: C, 79.47; H, 7.07; N, 3.50.

REFERENCES

1. R. B. Nielsen, *J. Phys. Chem.*, **68**, 2012 (1964).
2. R. Quast and R. Bieder, *Angew. Chem.*, **85**, 415 (1973).
3. R. B. Nielsen and J. G. Clifton, *J. Exp. Chem.*, **49**, 314 (1975).
4. R. Quast, E. Scheller, and E. Frank, *Angew. Chem. International Ed.*, **1**, 451 (1972).
5. E. B. Pollard and E. F. Bertozzi, *J. Amer. Chem. Soc.*, **73**, 3195 (1951).
6. A. T. Bellard and J. B. Boffroy, *J. Amer. Chem. Soc.*, **73**, 3445 (1951).
7. R. C. Weillinger and E. Kennedy, *Chem. Ind. (London)*, 348 (1965).
8. A. T. Bellard and E. L. Rice, *J. Amer. Chem. Soc.*, **83**, 191 (1961).
9. J. A. Seyrup and E. F. Bertozzi, *Techniques of Org. Chem.*, **3**, 309 (1963).
10. J. G. Clifton and E. B. Nielsen, *J. Amer. Chem. Soc.*, **93**, 4294 (1971).
11. E. F. Bertozzi and R. Bieder, *J. Amer. Chem. Soc.*, **89**, 3125 (1967).
12. W. Gesselt, T. Nischlich, and T. M. Kolligly, *J. Amer. Chem. Soc.*, **85**, 1330 (1963); L. B. Jellum and S. G. Schierblinger, *J. Exp. Chem.*, **78**, 1503 (1963); F. B. Jensen and W. E. Thomas, *J. Amer. Chem. Soc.*, **84**, 3348 (1962); P. Schuster, J. Schuster, S. Gendin, S. Gaffney, and G. Sefenich, *ibid.*, **84**, 3394 (1962); T. Nischlich, *J. Org. Chem.*, **26**, 7 (1961).
13. Spectral frequencies consistent with equation (2) at 1620 cm^{-1} , at 1530 cm^{-1} , $\nu_{\text{max}} = 134\text{ cm}^{-1}$, $123,490\text{ cm}^{-1}$.
14. Spectral data for cyclopentadienyl anion (2): Exact mass $\text{C}_5\text{H}_5\text{N} = 77.0439$, $\text{C}_5\text{H}_5\text{N} = 77.0439$, $\nu_{\text{max}} = 1620\text{ cm}^{-1}$, 1530 cm^{-1} , 134 cm^{-1} , $123,490\text{ cm}^{-1}$, 120 cm^{-1} , 110 cm^{-1} , 100 cm^{-1} , 90 cm^{-1} , 80 cm^{-1} , 70 cm^{-1} , 60 cm^{-1} , 50 cm^{-1} , 40 cm^{-1} , 30 cm^{-1} , 20 cm^{-1} , 10 cm^{-1} , 0 cm^{-1} .
15. J. G. Clifton and E. B. Nielsen, *J. Exp. Chem.*, **78**, 440 (1963).
16. R. B. Bieder and J. B. Boffroy, *J. Experimental Chem.*, **3**, 31 (1963).
17. E. Scheller and R. Quast, *Angew. Chem. Int. Ed.*, **4**, 1076 (1965).
18. J. A. Seyrup and E. F. Bertozzi, *J. Exp. Chem.*, **49**, 394 (1975).

11. J. Marchand-Brynaert and L. Dumas, *J. Amer. Chem. Soc.* **91**, 2887 (1969).
12. L. Hénaut, J. Marchand-Brynaert, and L. Dumas, *Angew. Chem. International Edition*, **11**, 180 (1972).
13. J. Marchand-Brynaert and L. Dumas, *J. Amer. Chem. Soc.* **91**, 2879 (1969).
14. R. De Pauwtere, J. Marchand-Brynaert, and L. Dumas, *Angew. Chem. International Edition*, **11**, 182 (1972).
15. L. Oestreich, E. Krensch, and R. C. Fieser, *Angew. Chem. International Edition*, **9**, 424 (1970); R. Fieser and L. Oestreich, *Translations Ann.* **196** (1972), reference 16.
16. C. L. Koenig, *Angew. Chem. International Edition*, **10**, 435 (1971).
17. L. Langryl and J. E. Hershman, *Angew. Chem. International Edition*, **3**, 31 (1964); J. E. Hershman and J. E. Hershman, *J. Amer. Chem. Soc.* **86**, 340 (1964); J. E. Hershman and L. Langryl, *J. Amer. Chem. Soc.* **86**, 116 (1964).
18. M. L. Huggins, J. Schell, and F. Fehsenfeld, *J. Amer. Chem. Soc.* **91**, 1909 (1969).
19. R. S. Stearns and R. Arceff, *J. Org. Chem.* **15**, 345 (1950).
20. R. Hershman and F. Fehsenfeld, *Angew. Chem. International Edition*, **3**, 388 (1964).
21. C. L. Koenig and J. G. French, *J. Amer. Chem. Soc.* **86**, 417 (1964).
22. C. L. Koenig and J. G. French, *J. Amer. Chem. Soc.* **86**, 4182 (1964).
23. C. L. Koenig and E. R. Kligel, *J. Amer. Chem. Soc.* **86**, 54 (1964).
24. R. J. Fieser, J. Linnert, and L. Rott, *Angew. Lichtst. Int. Chem.* **7**, 79, 84 (1966).
25. K. M. Lee and E. A. Slinger, *J. Org. Chem.* **29**, 1158 (1964).
26. R. C. Cook and W. C. Fernelius, *J. Amer. Chem. Soc.* **86**, 944 (1964).
27. This material although not isolated exhibited a medium intensity infrared absorption band at 2160 cm^{-1} .
28. L. J. Bellamy, "The Infrared Spectra of Complex Molecules," 2nd ed., John Wiley and Sons, Inc., New York, 1958, (212).
29. R. H. Williams and L. Fleming, "Spectroscopic Methods in Organic Chemistry," 2nd ed., McGraw-Hill, New York, 1972.

19. F. Harper and L. Benaghi-Labib, *Chem. Reviews*, **42**, 567 (1961).
20. Methyl Chlorosulfonate (Mega Ketyl), 97% was purchased from Aldrich Chemical Co., Inc. This reagent has been established to be a reagent grade. All manipulations were carried out in a well ventilated hood and protective rubber gloves were also being worn.
21. R. Garfield, W. J. Walp, and J. T. Herold, *J. Amer. Chem. Soc.*, **78**, 1286 (1956).
22. F. A. Rasey, "Radical Organic Synthesis Spectroscopy," Academic Press, New York, 1961, p. 107.
23. T. Kato-Mae and C. H. Towns, *J. Amer. Chem. Soc.*, **78**, 1608 (1956).
24. R. Salinger, W. Schner, C. Tschelch, and R. Nator, *Zeitschrift für Naturforsch.*, **18b** (1963); R. Salinger, W. Schner, and R. Nator, *Angew. Chem. International Edn.*, **2**, 400 (1963); R. Salinger, W. Schner, R. Nator, and E. Korte, *Zeits.*, **18**, 426 (1963); R. N. Nator and R. Nator, *Zeitschrift für Naturforsch.*, **18b** (1963); R. N. Nator, R. N. Nator, E. J. Salinger, and R. J. Schuster, *J. Org. Chem.*, **28**, 2358 (1963).
25. R. Salinger, W. Schner, and R. Nator, *J. Amer. Chem. Soc.*, **78**, 2710 (1956); R. Salinger and R. Nator, *Angew. Chem. International Edn.*, **2**, 406 (1963).
26. R. N. Nator, W. J. Smith, and J. H. Nator, *J. Org. Chem.*, **25**, 1897 (1960); R. N. Nator and E. J. Nator, *Zeits.*, **15**, 171 (1960); R. N. Nator and E. Nator, *Zeits.*, **16**, 1461 (1961); E. Nator and R. Nator, *Zeits.*, **16**, 1461 (1961); E. Nator, *J. Org. Chem.*, **26**, 1201 (1961); E. Nator, *Zeits.*, **16**, 1461 (1961).
27. A. Fickel, J. Schuster, and R. L. Nator, Jr., *J. Org. Chem.*, **28**, 1591 (1963); R. Salinger, E. Nator, E. Nator, and E. Nator, *Chem. Rev.*, **43**, 1268 (1963).
28. G. C. Levy and C. L. Salinger, "Carbon-13 Radical Magnetic Resonance for Organic Chemistry," John Wiley and Sons, Inc., New York, (1961).
29. The empirical relationship $\log_{10} \nu_{\text{max}}^{13\text{C}} \approx 17.5 - 0.5 \log_{10} \nu_{\text{max}}^{12\text{C}}$ of the spectral signal or half height was used to estimate the magnitude of multiple coupling constants. E. Nator and E. J. Nator, *J. Org. Chem.*, **28**, 1591 (1963); R. N. Nator, R. N. Nator, R. J. Schuster, and R. N. Nator, *J. Amer. Chem. Soc.*, **78**, 4884 (1956).
30. J. J. Schuster and E. J. Nator, "The Nator's Transition," John Wiley and Sons, Inc., New York, 1970.
31. R. Nator and J. J. Schuster, *J. Amer. Chem. Soc.*, **78**, 4404 (1956); R. N. Nator and J. J. Nator, *J. Amer. Chem. Soc.*, **78**, 4766 (1956).

79. The 555 labeled compounds were prepared by coupling to the 555-labeled fluorescent yellow polyaniline derivative 53 which formed in the course of the same synthesis. Both 51 and 52 showed identical fragmentation patterns.
80. R. E. Miller, P. E. Brown, W. E. F. Mueller, and R. A. Frieseman, Chem. Commun., 143 (1964).
81. R. Bologan, Prog. Chem. Sci., 367 (1962).
82. R. Bologan, R. Gaudry, P. Lema, and R. Levesque, Angew. Chem., 75, 418 (1963).
83. R. Bologan, W. Schanz, and G. Gauthier, Integrations (1962), 367 (1962).
84. R. Bologan and P. Goss, Integrations (1962), 691 (1962).
85. R. E. Peubeard and L. Hoffmann, "The Conservation of Orbital Symmetry," Academic Press, New York, 1965.
86. R. J. Hirst, R. E. Pease, and R. T. Gortan, "The Systematic Identification of Organic Compounds," 3th ed., John Wiley and Sons, Inc., New York, 1962, pp. 115-161.
87. E. S. Lewis, L. F. Long, and H. Ye Lohian, Ann. Repts. Chem. Soc., 574 (1961).
88. R. E. Rether and J. R. Bennett, J. Macromol. Chem., 5, 1487 (1961).
89. R. E. Rether and J. R. Bennett, J. Macromol. Chem., 5, 1497 (1961).
90. R. E. Rether and J. R. Bennett, J. Macromol. Chem., 11, 361 (1964).
91. R. E. Rether, "Infrared Spectroscopy," McGraw and Hill, Inc., Boston, 1964, p. 163.
92. C. J. Forehand and J. C. Easfield, "The Aldrich Library of IR Spectra," Vol. VI, Aldrich Chem. Co., Inc., 1959.
93. L. A. Singer, L. A. Davis, and R. L. Benson, J. Amer. Chem. Soc., 75, 1148 (1953).
94. R. J. Leonard and F. W. Smith, J. Amer. Chem. Soc., 82, 5192 (1960).
95. G. A. Olah and S. Trachtenberg, J. Amer. Chem. Soc., 89, 4716 (1967).
96. "Kautlin Index," Kautlin Research Laboratories, 1018 Spring Garden Street, Philadelphia, Pa., (1964).
97. G. S. Farnar and J. Goldstein, "Mass Spectrometry of Macromolecular Compounds," John Wiley and Sons, Inc., New York, 1963, pp. 188-346.
98. R. E. Hill, Jr. and P. Thoms, J. Amer. Chem. Soc., 91, 606 (1969).

11. E. Wehrha, F. Knechtig, and R. Koser, Angew. Math. Physik, **20**, 247 (1971).
12. E. Loring and G. Baker, J. Chem. Phys., **23** (1955).
13. R. Velupillai, J. Org. Chem., **35**, 3478 (1970).
14. E. Dineen, E. Bauer, and J. Smith, Analyst. Chem., **43**, 241 (1971).
15. R. A. S. Smith and T. J. Tennil, J. Org. Chem., **35**, 1475 (1970).
16. J. March, "Advanced Organic Chemistry: Reactions, Mechanisms, and Synthesis," (McGraw-Hill, Inc., New York, 1967), pp. 274-187.
17. L. Fieser, "Principles of Modern Heterocyclic Chemistry," (Wiley, New York, 1968), p. 108; G. S. Hammond, J. Amer. Chem. Soc., **73**, 543 (1951); R. W. Hurley and R. S. Kaplan, J. Amer. Chem. Soc., **85**, 577 (1963).
18. J. A. Dwyer, Introduction, **1971** (1971).
19. J. A. Dwyer, J. Org. Chem., **35**, 2115 (1970).
20. Initial reaction times of 20-40 hours produced maximum $\Delta\epsilon$ but no lower yields.
21. P. E. Miller and E. I. Gormell, J. International. Chem., **3**, 179 (1969).
22. N. Toullec, R. Sauter-Breger, E. Gaudin, J. Baudouin, and E. Carrière, Introduction, **1971** (1971).
23. P. Toullec and E. Carrière, Introduction, **1971** (1971).
24. J. C. Gild, Ph.D. Thesis, University of Florida, 1971.
25. E. T. Yen, Masters Thesis, University of Florida, 1970.
26. J. A. Dwyer, R. W. Neill, W. E. Pogue, and E. T. Yen, Introduction, **1971** (1971).
27. E. S. Mayerson and R. S. Scheraga, J. Amer. Chem. Soc., **85**, 1566 (1963).
28. R. Gilman, P. S. Williams, M. P. Fisher, and G. R. Kuper, J. Amer. Chem. Soc., **85**, 145 (1963).
29. R. S. Stein and P. W. Smith, "Handbook of Chemistry and Physics," 41st ed., (The Chemical Rubber Co., Cleveland, Ohio, 1968-69).
30. R. Steiniger, Int. J. Radiat. Chem. Phys., **15**, 544 (1971).
31. R. Steiniger, Angew. Math. Phys. Phys., **20**, 81 (1969).

11. R. Knechtger and E. Knecht, *Helv. Chim. Acta*, **3**, 517 (1921).
12. R. G. Catwood, Jr. and J. C. Goff, *J. Amer. Chem. Soc.*, **72**, 3117 (1950).
13. R. L. Naffziger, R. L. Sargent, and R. E. Spencer, *J. Amer. Chem. Soc.*, **77**, 5437 (1955).
14. R. S. Steinert and J. F. Jenkins, Jr., *J. Amer. Chem. Soc.*, **75**, 3347 (1953).
15. R. Knapton and R. M. Brown, *J. Amer. Chem. Soc.*, **77**, 3353 (1955).
16. T. Rostrom, *Industria Saccar.*, **32**, 415 (1945).
17. A. P. Frankhamer, *Proc. 2. Intern. Symposium*, **35**, 395 (1949).
18. R. Fischer, *Angew. Chem. Intern. Ed.*, **4**, 412 (1965).
19. L. Sri, L. Meyer, W. Lichinski, F. Schindler, and E. Kuntzsch, *Helv. Chim. Acta*, **33**, 13 (1950).
20. A. Forni-Rossini, L. M. Kryukova, and L. Tie. Sella, *Chim. Ind. (Milan)*, **35**, 1073 (1953).
21. W. Sella and G. Scherzoff, *Helv. Chim. Acta*, **33**, 140 (1950); C. A. Hickhoff and A. Broadwater, *Angew. Chem. Intern. Ed.*, **10**, 3179 (1961).

THE BUREAU






Fig. 1. Spectrum 1.




THE GARDEN



WIR SPEKTREN 1



FIGURE 3
 1000 2000 3000 4000 5000 6000 7000 8000 9000 10000






1. *Salix caprea* L. (Weeping Willow)
 2. *Salix viminalis* L. (Common Willow)
 3. *Salix purpurea* L. (Purple Willow)
 4. *Salix alba* L. (White Willow)
 5. *Salix pyramidalis* B. (Pyramidal Willow)
 6. *Salix triandra* L. (Triandra Willow)
 7. *Salix glauca* L. (Blue Willow)
 8. *Salix elaeagnifolia* L. (Elaeagnifolia Willow)
 9. *Salix virens* L. (Virens Willow)
 10. *Salix caprea* L. (Weeping Willow)


Willow (Salix)

DATE: / /






NMR SPECTRUM 6




1H NMR spectrum of 1,1-dimethyl-4-propylcyclohexane




Sketch of a rock


MS SPECTRUM 11




100% CDCl_3



MAN SPECTRUM 13



OAS SPECTRUM 14







FIG. 1. SPECTRUM 13

100% SPECTRUM 1A



208. SPECTRUM 12





Fig. 1. Spectrum of 1.



12. KOMPONENTE 12








FIG. SECTION 3.1




12. NUTRITION 21



van der Vliet et al.





COSY SPECTRUM 12



UNEMPLOYED STATE

George Hughes Randolph Egan was born March 15, 1917, in Northampton, Massachusetts. Shortly thereafter, his family moved to Washington, D.C. where in June, 1935, he was graduated from George College High School. In June, 1938, he received a Bachelor of Science degree with a major in Chemistry from Rollins College, Winter Park, Florida. During the following year, he was employed as an analytical chemist by Rivers Chemical Laboratories, Altamonte Springs, Florida. In September, 1939, Mr. Egan enrolled as a graduate student at Wake Forest University, Winston-Salem, North Carolina where in August, 1941, he received a Master of Arts degree with a major in Inorganic Chemistry. In September, 1943, he enrolled in the Graduate School of the University of Florida to pursue doctoral studies in the Department of Chemistry. During his graduate study, Mr. Egan held a teaching assistantship from September, 1943 to December, 1945, and, in 1946, received the Robert Award for Excellence in Teaching. Upon graduation, he will begin postdoctoral study at the University of Rochester with Professor Arthur Ewing.

I certify that I have read this study and that in my opinion it conforms to acceptable standards of scholarly presentation and is fully adequate, in scope and quality, as a dissertation for the degree of Doctor of Philosophy.


James A. Mayrow
Professor of Chemistry

I certify that I have read this study and that in my opinion it conforms to acceptable standards of scholarly presentation and is fully adequate, in scope and quality, as a dissertation for the degree of Doctor of Philosophy.


William H. Jones
Professor of Chemistry

I certify that I have read this study and that in my opinion it conforms to acceptable standards of scholarly presentation and is fully adequate, in scope and quality, as a dissertation for the degree of Doctor of Philosophy.


William S. Jenkins, Jr.
Professor of Chemistry

I certify that I have read this study and that in my opinion it conforms to acceptable standards of scholarly presentation and is fully adequate, in scope and quality, as a dissertation for the degree of Doctor of Philosophy.


Gus J. Peltz
Professor of Chemistry

I certify that I have read this study and that in my opinion it conforms to acceptable standards of scholarly presentation and is fully adequate, in scope and quality, as a dissertation for the degree of Doctor of Philosophy.



Earl M. Dean
Assistant Professor of
Biochemistry

This dissertation was submitted to the Graduate Faculty of the Department of Chemistry in the College of Arts and Sciences and to the Graduate Council, and was accepted as partial fulfillment of the requirements for the degree of Doctor of Philosophy.

December, 1956

Dean, Graduate School